

1 **Enhancing life cycle impact assessment from climate science: Review of**
2 **recent findings and recommendations for application to LCA**

3 Annie Levasseur^{*a}, Otávio Cavalett^b, Jan S. Fuglestedt^c, Thomas Gasser^{d,e}, Daniel J.A.
4 Johansson^f, Susanne V. Jørgensen^g, Marco Rauegi^h, Andy Reisingerⁱ, Greg Schivley^j, Anders
5 Strømman^k, Katsumasa Tanaka^l, Francesco Cherubini^k

6 Corresponding author e-mail: annie.levasseur@polymtl.ca

7 Corresponding author phone: +1-514-340-4711 #4013

8 ^a CIRAIG, Department of Chemical Engineering, Polytechnique Montréal, P.O. Box 6079, Stn
9 Centre-ville, Montréal, Québec, H3C 3A7, Canada, annie.levasseur@polymtl.ca

10 ^b Laboratório Nacional de Ciência e Tecnologia do Bioetanol (CTBE), Centro Nacional de
11 Pesquisa em Energia e Materiais (CNPEM), Caixa Postal 6192, CEP 13083-970, Campinas, São
12 Paulo, Brasil, otavio.cavalett@bioetanol.org.br

13 ^c Center for International Climate and Environmental Research – Oslo (CICERO), Oslo, Norway,
14 j.s.fuglestedt@cicero.oslo.no

15 ^d Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Institut Pierre-Simon
16 Laplace (IPSL), CEA-CNRS-UVSQ, CEA l'Orme des Merisiers, 91191 Gif-sur-Yvette Cedex,
17 France, thomas.gasser.2006@polytechnique.org

18 ^e Centre International de Recherche sur l'Environnement et le Développement (CIRED), CNRS-
19 PontsParisTech-EHESS-AgroParisTech-CIRAD, Campus du Jardin Tropical, 45bis avenue de la
20 Belle Gabrielle, 94736 Nogent-sur-Marne Cedex, France

21 ^f Division of Physical Resource Theory, Department of Energy and Environment, Chalmers
22 University of Technology, Gothenburg, Sweden, daniel.johansson@chalmers.se

23 ^g ALECTIA A/S, Teknikerbyen 34, 2830 Virum, Denmark, sus_vj@hotmail.com

24 ^h Faculty of Technology, Design and Environment, Oxford Brookes University, Wheatley, UK,
25 marco.rauegi@brookes.ac.uk

26 ⁱ New Zealand Agricultural Greenhouse Gas Research Centre, Private Bag 11008, Palmerston
27 North 4442, New Zealand, andy.reisinger@nzagrc.org.nz

28 ^j Civil and Environmental Engineering, Carnegie Mellon University, Pittsburgh, Pennsylvania,
29 United States, gs1@cmu.edu

30 ^k Industrial Ecology Programme, Department of Energy and Process Engineering, Norwegian
31 University of Science and Technology (NTNU), Trondheim, Norway,
32 anders.hammer.stromman@ntnu.no, francesco.cherubini@ntnu.no

33 ⁱ National Institute for Environmental Studies (NIES), Tsukuba, Japan,
34 tanaka.katsumasa@nies.go.jp

35

36 **Abstract.**

37 Since the Global Warming Potential (GWP) was first presented in the Intergovernmental Panel on
38 Climate Change (IPCC) First Assessment Report, the metric has been scrutinized and alternative
39 metrics have been suggested. The IPCC Fifth Assessment Report gives a scientific assessment of
40 the main recent findings from climate metrics research and provides the most up-to-date values
41 for a subset of metrics and time horizons. The objectives of this paper are to perform a systematic
42 review of available midpoint metrics (i.e. using an indicator situated in the middle of the cause-
43 effect chain from emissions to climate change) for well-mixed greenhouse gases and near-term
44 climate forcers based on the current literature, to provide recommendations for the development
45 and use of characterization factors for climate change in life cycle assessment (LCA), and to
46 identify research needs. This work is part of the ‘Global Guidance on Environmental Life Cycle
47 Impact Assessment’ project held by the UNEP/SETAC Life Cycle Initiative and is intended to
48 support a consensus finding workshop. In an LCA context, it can make sense to use several
49 complementary metrics that serve different purposes, and from there get an understanding about
50 the robustness of the LCA study to different perspectives and metrics. We propose a step-by-step
51 approach to test the sensitivity of LCA results to different modelling choices and provide
52 recommendations for specific issues such as the consideration of climate-carbon feedbacks and
53 the inclusion of pollutants with cooling effects (negative metric values).

54 **Keywords.** Climate change, Life cycle assessment (LCA), Climate metric, Well-mixed
55 greenhouse gas, Near-term climate forcer

56 **1. Introduction**

57 Life cycle assessment (LCA) is a decision support tool that estimates the potential environmental
58 impacts of any product system over its entire life cycle. It is commonly used to guide
59 environmental policies and programs, to inform consumers' choices through environmental
60 labeling and declarations, and to help industries reduce the environmental impact of their
61 activities or design more sustainable products, amongst others (ISO 14044, 2006).

62 The first step in an LCA – after defining the goal and scope – is to develop an inventory of all
63 environmental emissions from, and natural resource inputs to, each unit process in the system.
64 The total environmental inputs and outputs from all activities are called the life cycle inventory
65 (LCI). In life cycle impact assessment (LCIA), these environmental flows are classified according
66 to the type of environmental impact they cause, and multiplied by characterization factors (CF)
67 that express their contribution to that indicator. CFs are developed using environmental models
68 that estimate the relative or absolute effect of each flow on a selected indicator, which is a
69 quantifiable representation of an impact category. LCA practitioners usually select a specific
70 LCIA method that proposes a series of CFs for different types of environmental impact (ISO
71 14044, 2006).

72 Emissions of CO₂ and other greenhouse gases (GHGs), aerosols, and ozone precursors are
73 affecting the climate system as illustrated by the cause-effect chain presented in Figure 1. In
74 current LCIA methods, CFs for the climate change impact category are usually proposed only for
75 well-mixed greenhouse gases (WMGHG), using Global Warming Potential (GWP) values
76 published in the Intergovernmental Panel on Climate Change (IPCC) assessment reports. Other
77 anthropogenic causes of global warming such as near-term climate forcers (NTCF) or albedo
78 changes are currently not considered in LCA (Levasseur, 2015). The important difference
79 between WMGHGs and NTCFs is their lifetime. WMGHGs have atmospheric lifetimes long

80 enough to be well mixed throughout the troposphere, and their climatic impact does not depend
81 on the location of emissions. WMGHGs include CO₂, N₂O, CH₄, SF₆ and many halogenated
82 species. By contrast, NTCFs have atmospheric lifetimes of less than one year so that their
83 climatic impact depends on the emission location. NTCFs include ozone and aerosols, or their
84 precursors, and some halogenated species that are not WMGHGs (Myhre et al., 2013).

85 Researchers have shown that cumulative emissions of WMGHG with a lifetime greater than 50-
86 100 years dominate the peak warming (Smith et al., 2012). However, reducing emissions of
87 NTCFs and WMGHGs with shorter lifetimes could reduce the rate of climate warming over the
88 next few decades and, if emission reductions are sustained, also lower the peak temperature
89 attained (Myhre et al., 2011; Penner et al., 2010; Rogelj et al., 2014; Shindell et al., 2012; Smith
90 et al., 2012). If net CO₂ emissions do not decline significantly and eventually reach zero,
91 mitigation of short-lived species will only postpone but not avoid the breaching of a temperature
92 threshold in line with those adopted within the UNFCCC process (Allen et al., 2016; Bowerman et
93 al., 2013).

94 There are two different types of CFs depending on the position of the selected indicator in the
95 cause-effect chain (see Figure 1). Midpoint CFs refer to effects at an earlier stage of the cause-
96 effect chain such as radiative forcing or temperature, while endpoint CFs are derived from
97 relatively more complex mechanisms (with increased uncertainties) for translating emissions into
98 impacts on human health (e.g. disability-adjusted life years caused by climate change) and
99 ecosystems (e.g. potential disappeared fraction of species because of climate change) (Levasseur,
100 2015). All current LCIA methods offer midpoint CFs using GWPs published by the IPCC. The
101 only distinctions between LCIA methods on this matter are the choice of time horizon and the
102 issue year of the IPCC Assessment Report. Most LCIA methods use a 100-year time horizon to
103 be in line with the time horizon selected for the application in the 1997 Kyoto Protocol, while a
104 very few others use a 20- or 500-year time horizon (Levasseur, 2015). For instance, the ReCiPe

105 method uses time horizons of 20, 100 and 500 years respectively for the individualist, hierarchist
106 and egalitarian perspectives (Goedkoop et al., 2013). Users must choose between one of these
107 perspectives to set the default value for some modeling choices.

108 Since the GWP was first presented in the IPCC First Assessment Report, the metric has been
109 scrutinized and alternative metrics have been suggested. GWP was intended to clarify the relative
110 contributions to global warming of different countries and different activities to help develop
111 cost-effective emission policies at both national and international levels (Lahosf & Ahuja, 1990).
112 However, Shine (2009) reminds us that the GWP concept was initially a simple approach adopted
113 in part to illustrate the difficulties encountered when developing a single metric to assess climate
114 impacts associated with GHG emissions of gases with very different physical and chemical
115 properties. There exists a plethora of other metrics based on physical and biogeochemical aspects
116 of climate change (e.g. Gillet & Matthews, 2010; Lauder et al., 2013; Peters et al., 2011a; Shine
117 et al., 2005, 2015; Smith et al., 2012; Sterner et al., 2014; Tanaka et al., 2009; Wigley, 1998), and
118 a large range of metrics where aspects of economics are also taken into account (e.g. Eckaus,
119 1992; Johansson, 2012; Manne & Richels, 2001; Rillee & Richards, 1993). In recent years, the
120 issue of metrics has received increased political attention and several publications have addressed
121 concerns regarding the use of appropriate climate metrics in an LCA context (e.g. Peters et al.,
122 2011b; UNFCCC, 2012, 2014). The IPCC Fifth Assessment Report (5thAR) gives a scientific
123 assessment of the main recent findings from physical climate metrics research and provides the
124 most up-to-date values for a subset of metrics and time horizons (GWP and Global Temperature
125 change Potential (GTP); see below). Crucially, the latest IPCC assessment emphasises that the
126 choice of emission metric and time horizon depends on type of application and policy context and
127 no single metric is optimal for all policy goals (IPCC, 2014a).

128 The objectives of this paper are to perform a systematic review of available midpoint metrics for
129 WMGHGs and NTCFs based on the current literature, to provide recommendations for the

130 development and use of climate change CFs in LCA, and to identify research needs. We primarily
131 discuss research findings on metrics presented in the IPCC 5thAR, which emphasized GWP and
132 GTP, and under which circumstances these metrics could be applied to improve current climate
133 change midpoint characterization factors in LCA. This work is part of the ‘Global Guidance on
134 Environmental Life Cycle Impact Assessment’ project held by the UNEP/SETAC Life Cycle
135 Initiative and is intended to support a consensus finding workshop

136 **2. Emission metrics for climate change impacts**

137 Emission metrics aim to compare the effects of different forcing agents on the climate system.
138 They can be used in different contexts such as multi-component climate policies, comparison of
139 emissions between regions or sectors, and LCA, amongst others (Kolstad et al., 2014; Myhre et
140 al, 2013). As stated in the IPCC 5thAR, “the most appropriate metric will depend on which
141 aspects of climate change are most important” (Myhre et al, 2013). Indeed, no single metric can
142 adequately and simultaneously assess the impact of different climate forcers on different aspects
143 of climate change such as the rate of change or long-term temperature increase. This section
144 presents an overview of different midpoint emission metrics used to estimate the impact of
145 climate forcers.

146 **2.1. Development of emission metrics**

147 Metrics are intended to be applied widely and with minimized value-judgements, but a number of
148 choices have to be made in order to select a specific metric (Tanaka et al., 2010). This section
149 presents the cause-effect chain of climate change from which an indicator must first be selected,
150 as well as fundamental choices about some metric characteristics.

151

152


153

2.1.1. The cause-effect chain

154 The emission of WMGHGs and NTCFs leads to an increase of their concentration in the
155 atmosphere. This increase in concentration results in radiative forcing [$\text{W}\cdot\text{m}^{-2}$], also called
156 stratospherically adjusted radiative forcing, which is defined as the change in net downward
157 radiative flux at the tropopause after the stratospheric temperature is readjusted to a radiative
158 equilibrium, while surface and tropospheric temperature and state variables such as water vapour
159 and cloud cover are held fixed at the unperturbed values (Myhre et al., 2013). An extension to
160 this, the effective radiative forcing, considers the change in radiative flux after allowing for an
161 adjustment of other physical variables such as cloud and snow cover. Effective radiative forcing
162 is nearly identical to radiative forcing for WMGHGs but can differ significantly for aerosols
163 because of their additional direct impact on clouds and snow cover. A positive radiative forcing
164 warms the climate system, while a negative radiative forcing cools it.

165 WMGHGs and NTCFs cause radiative forcing when they are released in the atmosphere, which
166 thus leads to a warming (most WMGHGs and NTCFs) or cooling (some NTCFs) impact on the
167 land and sea surface temperature. This temperature change leads to different climate impacts such
168 as sea-level rise, changes in precipitations, melting of polar ice-caps and glaciers, thawing of
169 permafrost, etc. Aerosols can also affect climate directly through influencing cloud properties and
170 formation and snow melt. These changes finally impact humans and ecosystems in different ways
171 such as by increasing the incidence of certain diseases, flooded areas, changes in food production,
172 droughts leading to malnutrition, species range shifts and possible extinctions etc. (IPCC, 2014b).
173 Figure 1 presents a simplified diagram of the cause-effect chain from emissions to impacts as
174 described above. The diagram shows only the most important pathways by omitting more detailed
175 feedbacks such as climate-carbon cycle feedbacks (Friedlingstein et al., 2006).

176 Emission metrics are often used to quantify and compare the climate impacts of WMGHGs and
 177 NTCFs in different accounting methodologies such as LCA or GHG emission inventories. They
 178 are developed by choosing an indicator somewhere in the cause-effect chain and measuring the
 179 effect of an emission of each climate forcer on this indicator based on input from climate
 180 modelling. This paper focuses on midpoint indicators only.



181

182 Figure 1. The cause-effect chain of climate change (modified from Figure 8.27 of IPCC 5th AR
 183 WGI)

184 **2.1.2. Absolute versus normalized metrics**

185 Absolute metrics estimate the value of the selected indicator for a given emission (e.g. Kelvin per
 186 unit emitted), while normalized metrics compare the value of the selected indicator for a given
 187 emission to the value of the same indicator for an equal mass of a reference substance emitted

188 (i.e. a ratio of an absolute metric of a component of interest to the absolute metric of a reference
189 gas). The reference substance used conventionally for climate metrics is CO₂, with emissions
190 commonly expressed as so-called “CO₂-equivalent” emissions. While this may be justified given
191 that CO₂ is the dominant cause of human-induced warming, this choice also introduces
192 complexity related to the variety of physical and biogeochemical processes governing the
193 atmospheric CO₂ concentration (namely the non-single exponential decay of a pulse emission
194 into the atmosphere, with a fraction remaining for many thousands of years) as well as the fact
195 that the climate effect due to a normalised pulse emission of this gas itself changes over time with
196 changing background concentrations and on-going climate change (Joos et al., 2013).

197 **2.1.3. Instantaneous versus cumulative metrics**

198 For the majority of suggested metrics it is possible to categorize them as either being
199 instantaneous or cumulative. Instantaneous metrics estimate the value of the selected indicator at
200 a given point in time after an emission (“snapshot metrics”), while cumulative metrics integrate
201 the value of the selected indicator over a period of time up to a given time horizon. Given the
202 very different atmospheric lifetimes of different GHGs, the choice between instantaneous and
203 cumulative metrics has a fundamental bearing on the value accorded to relatively short-lived
204 gases relative to those with longer lifetimes, as cumulative metrics ‘remember’ near-term
205 warming even if a distant time horizon is chosen for evaluating climate impacts (Fuglestedt et
206 al., 2010; Myhre et al, 2013; Sterner et al., 2014).

207 **2.2. Presentation of selected climate metrics**

208 This section presents some specific instantaneous and cumulative metrics based on different
209 indicators such as radiative forcing and temperature change that have robust literature behind
210 them, some of them having been assessed by the IPCC. Each of these metrics embodies

211 implications of different choices at various points in their development. Some of these choices are
212 discussed further in Section 3.

213 **2.2.1. Metrics based on radiative forcing**

214 Some metrics use the radiative forcing caused by an emission as key parameter for indicating
215 impacts of different climate forcers. Different models and assumptions can be used to estimate
216 radiative forcing, which depends on the radiative efficiency of the climate forcer, its atmospheric
217 lifetime by various removal processes, and any indirect effects.

218 The most widely used metric based on radiative forcing is GWP. This is largely because it was
219 the only option considered by the IPCC in its First Assessment Report (IPCC, 1990) and adopted
220 for the application of the Kyoto Protocol. GWP [$\text{kgCO}_2\text{-eq}\cdot\text{kg}^{-1}$] is an example of a normalized
221 and cumulative metric. It is the ratio of the Absolute Global Warming Potential (AGWP) of a
222 given GHG to that of CO_2 , the reference gas (Equation 4). AGWP [$\text{W}\cdot\text{yr}\cdot\text{m}^{-2}\cdot\text{kg}^{-1}$], an absolute
223 metric, is the cumulative radiative forcing caused by a unit-mass pulse emission calculated over a
224 selected time horizon TH [yr] (Equation 1) (Myhre et al., 2013).

$$225 \quad \text{AGWP}_x(\text{TH}) = \int_0^{\text{TH}} \text{RF}_x(t) dt \quad (1)$$

226 For non- CO_2 gases, $\text{RF}(t)$ can be approximated by a first-order decay equation (see Equation 2)
227 where A [$\text{W}\cdot\text{m}^{-2}\cdot\text{kg}^{-1}$] is the radiative efficiency i.e. the radiative forcing caused by a marginal
228 increase in atmospheric concentration of a given gas in the atmosphere and τ [yr] is the lifetime of
229 the gas. Additional terms or adjustments may be needed in Equations 1 and 2 to account for
230 effects such as climate-carbon cycle feedbacks or the oxidation of fossil CH_4 to CO_2 .

$$231 \quad \text{RF}_x(t) = A_x e^{-t/\tau_x} \quad (2)$$

232 For CO₂, in IPCC 5th AR as well as previous reports, $RF(t)$ is given by a more complex formula
233 (see Equation 3) derived as the average response across multiple carbon-cycle models of various
234 complexities (Joos et al., 2013). This represents the response of the oceanic and terrestrial carbon
235 sinks to an instantaneous increase in atmospheric CO₂.

$$236 \quad RF_{CO_2}(t) = A_{CO_2} \left\{ a_0 + \sum_{i=1}^3 a_i e^{-t/\tau_i} \right\} \quad (3)$$

$$237 \quad GWP_x(TH) = \frac{AGWP_x(TH)}{AGWP_{CO_2}(TH)} \quad (4)$$

238 In each of its assessment reports, the IPCC publishes a list of updated values for the different
239 parameters A , a , and τ [yr] for a constant (but updated) atmospheric background concentration, as
240 well as GWP values calculated for some selected time horizons. Traditionally, GWP values were
241 calculated for 20-, 100-, and 500-year time horizons. However, in its 5th AR, the IPCC shows
242 values for 20- and 100-year time horizons only, stating that the confidence in providing useful
243 metrics for time horizons longer than 100 years is very low due to associated uncertainties and
244 strong assumptions of constant background conditions as well as ambiguity in the interpretation
245 of such a long-integration metric (Myhre et al., 2013).

246 AGWP and GWP are cumulative metrics. However, a metric based on instantaneous radiative
247 forcing using the same parameters as those used for AGWP could be developed using the value of
248 $RF(t)$ at a given time instead of integrating it over the time horizon (Edwards & Trancik, 2014;
249 Michaelis, 1992). Such a metric would estimate the change in radiative forcing occurring at any
250 given time following an emission. Another example of forcing-based metric is the Forcing
251 Equivalence Index (FEI), a time-dependent metric to capture climate forcing along a prescribed
252 scenario (Manning & Reisinger, 2011; Wigley, 1998).

253

254

255 **2.2.2. Metrics based on temperature change**

256 Some metrics go one-step further down the cause-effect chain and use the temperature change
257 caused by an emission as an indicator to assess the impact of different climate forcings. The Global
258 Temperature change Potential (GTP) proposed by Shine et al. (2005) is a normalized metric and
259 was assessed in the IPCC 5thAR. It is an example of an instantaneous metric, defined as the
260 change in global mean surface temperature at a chosen point in time TH [yr] after a pulse-
261 emission, relative to the temperature change following a pulse emission of a unit quantity of CO₂.
262 GTP [kgCO₂-eq·kg⁻¹] uses the same parameters as GWP i.e. radiative efficiency A [W·m⁻²·kg⁻¹]
263 and atmospheric decay τ [yr], as well as the climate sensitivity and the exchange of heat between
264 the atmosphere and the ocean using parameters c [K·(W·m⁻²)⁻¹] and d [yr]. Formulas have been
265 proposed for sustained and pulse emissions, but only the ones for pulse emissions are presented in
266 the IPCC 5th AR. The original AGTP formula has a single term for the time response of the
267 climate system (Shine et al, 2005). Since then, some researchers have proposed formulas with
268 two (Boucher & Reddy, 2008; Fuglestedt et al., 2010; Geoffroy et al., 2013) and three response
269 terms (Li & Jarvis, 2009; Olivie et al., 2012) to better represent the different time scales of the
270 climate response. However, Li and Jarvis (2009) argue that too many terms are difficult to
271 calibrate so that it is better to restrict to a two-term function. As an alternative to use analytical
272 impulse response functions to estimate metrics, one can use numerical reduced-complexity
273 climate models, such as MAGICC and ACC2 (e.g. Gillet & Matthews, 2010; Reisinger et al.,
274 2010, 2011; Tanaka et al, 2009, 2013).

275 Because metrics based on temperature change are midpoint indicators further down the cause-
276 effect chain than those using radiative forcing only as the key parameter, they may be more
277 relevant for determining environmental consequences of emissions, even if they use additional
278 uncertain parameters as indicated by Figure 1 in this paper as well as Figure 8.27 in Working

279 Group I IPCC 5th AR (Myhre et al, 2013). Indeed, the uncertainty in GWP and GTP cannot be
280 directly compared since they are of different nature. However, the GTP values also depend on the
281 response time of the climate system, which is uncertain. This uncertainty is a real feature of the
282 climate response which is not captured by the GWP (Fuglestvedt et al., 2003; Reisinger et al.,
283 2010).

284 Metrics based on cumulative temperature change over the selected time horizon, also known as
285 integrated GTP (iGTP), have been developed and analyzed (Azar & Johansson, 2012; Cherubini
286 et al., 2013; Gillet & Matthews, 2010; Peters et al., 2011a; Rotmans & den Elzen, 1992). The
287 iGTP for a given climate forcer and time horizon is under a range of circumstances approximately
288 similar to GWP for WMGHGs, but may be quite different for NTCFs such as black carbon.

289 TEMperature Proxy index (TEMP) is also a temperature-based metric (Tanaka et al., 2009). It is
290 defined for a given emission scenario and aims to capture the relative contribution of different
291 components to the temperature change.

292 Even though GTP is an instantaneous metric, it incorporates a degree of integration of radiative
293 forcing, since the temperature change at any given point in time reflects changes in radiative
294 forcing for up to several decades up to the temperature change. However, this ‘implicit
295 integration’ performed by the GTP is heavily weighted towards radiative forcing in the decade
296 immediately prior to the temperature change and depends on the time scales of the impulse
297 response function. This can also be seen by the fact that iGTP (rather than GTP) attains very
298 similar values to GWP for forcing agents with atmospheric lifetimes longer than a few years.
299 Thus the choice whether to adopt a cumulative or instantaneous metric is more important than the
300 implicit integration performed by metrics such as GTP which are further down the cause-effect
301 chain.

302

303 **2.2.3. Other types of metrics**

304 Sterner et al. (2014) have developed instantaneous and cumulative metrics to compare the impact
305 of different climate forcers on sea level rise: the Global Sea level rise Potential (GSP) and the
306 Integrated Global Seal level rise Potential (IGSP). They estimate the sea level rise at a given time
307 horizon (GSP) or the time integrated sea level rise (IGSP) caused by a pulse emission relative to
308 that of a comparable emission of CO₂. They have shown that all climate forcers, including very
309 short-lived ones, have considerable influence on sea level rise on the century time scale per unit
310 emissions.

311 Shine et al. (2015) present a new metric concept named the Global Precipitation change Potential
312 (GPP), which estimates the effect of various emissions on the global water cycle. The formulation
313 of GPP consists of two terms, one dependent on the surface temperature change and the other
314 dependent on the atmospheric component of the radiative forcing. For some forcing agents, and
315 notably for CO₂, these two terms oppose each other. Since the forcing and temperature
316 perturbations have different timescales, even the sign of the absolute GPP varies with time. One
317 finding is a strong near-term effect of CH₄ on precipitation change and the role of sustained
318 emissions of black carbon and sulphate in suppressing precipitation. The application of the GPP
319 in practice could be challenged by the fact that depending on location, an increase or decrease in
320 precipitation could be regarded as a negative or positive environmental impact.

321 The Climate Change Impact Potential (CCIP) is a metric that aims to capture a wide range of
322 climate impacts in an aggregated manner (Kirschbaum, 2014). While CCIP is by definition an
323 endpoint metric, it can be comparable to midpoint metrics because it is formulated as a function
324 of a set of three midpoint indicators: namely, instantaneous, cumulative, and rate of temperature
325 change. Quantifying climate impacts comprehensively is important but challenging, and this
326 metric inherently contains strong implicit assumptions. Most notable are i) an equal weighting


327 across the three impact terms using the midpoint indicators and ii) a fixed time horizon of 100
328 years.

329 Figure 2 presents an illustration of different kinds of metrics available and Table 1 shows the
330 values of metrics for non-fossil CH₄. The weighting of CH₄ relative to CO₂ varies substantially
331 between metrics and between time horizons, illustrating the significant implication of different
332 choices for climate metrics. Depending on the metric and time horizon chosen, the different
333 weight attributed to CH₄ emissions compared to CO₂ emissions could potentially lead to very
334 different conclusions when comparing product systems or climate mitigation solutions using
335 LCA.

336 Table 1. Values of midpoint metrics for non-fossil methane (kgCO₂-eq/kg) (FEI has been left out
337 of the table because its structure is different and it is not calculated for a fixed time horizon)

Metric	Time horizon = 20 years	Time horizon = 100 years
GWP (without feedbacks)	84	28
GWP (with feedbacks)	86	34
GTP (without feedbacks)	67	4
GTP (with feedbacks)	70	11
iGTP	81	28
GSP	78	18
IGSP	95	39
GPP	120	8.1
CCIP (using RCP6 scenario)	N/A	23

338



339

340 Figure 2. Illustration of different normalized metrics

341 **3. Discussion of some key metric choices**

342 This section discusses different key choices that one must make when selecting emission metrics.

343 These choices may have significant impacts on the LCA results. For instance, using GWP values

344 for a 20-year time horizon may lead to different conclusions than if GTP and a 100-year time

345 horizon is used. Despite the fact that science is able to inform decision makers about the

346 implications of these choices, it cannot objectively determine which ones are ultimately better

347 because it depends on policy context and involves value judgments so that there is no single

348 scientifically correct choice.

349

350

351 **3.1. Instantaneous versus cumulative metrics**

352 Climate impacts are diverse, and different types of emission metrics are needed to reflect the
353 different aspects, as summarized by Kirschbaum (2014). Instantaneous metrics based on
354 temperature are relevant when assessing the potential impacts caused by an absolute temperature
355 increase, which are closer to climate damages related for instance to heat waves (Huang et al.,
356 2011), extreme weather events (Webster et al., 2005), or coral bleaching (Baker et al., 2008).
357 Cumulative metrics based on temperature or radiative forcing are relevant when assessing
358 potential climate impacts associated with cumulative warming, which are for instance loss of
359 permanent ice and associated sea-level rise (Vermeer & Rahmstorf, 2009) or shut-off of
360 thermohaline circulation (Lenton et al., 2008). However, some researchers have shown that sea
361 level rise or thermohaline circulation, for instance, cannot be appropriately assessed by
362 cumulative metrics (Herrington & Zickfeld, 2014; Sterner et al., 2014), nor by instantaneous
363 metrics expressing the temperature increase in exactly one specific future year. Finally, the rate of
364 change of atmospheric and oceans temperature strongly influences whether species and humans
365 have time to adapt to climate change (Peck & Teisberg, 1994). A relevant metric for this type of
366 impacts would be based on the derivative of the function describing an instantaneous metric
367 (Hammit et al., 1996), but can also be reflected by the choice of shorter time horizons when using
368 a cumulative metric.

369 **3.2. Constant versus variable background atmosphere and climate**

370 GWP and GTP values proposed in the IPCC ARs have been calculated for respective present-day
371 constant background atmosphere concentrations (391 ppm CO₂ for the 5th AR) and climate
372 conditions. Values can also be calculated for variable conditions as already done (Joos et al.,
373 2013; Oliv  e et al., 2012; Reisinger et al., 2011; Tanaka et al., 2013). Considering variable
374 conditions may lead to more representative results since atmospheric concentrations and climate

375 conditions affect adjustment times and the concentration-forcing-temperature relationship (Myhre
376 et al., 2013). However, doing so requires assumptions to derive future emission scenarios such as
377 those used by the IPCC which increases the dependence of metric values on subjective
378 judgements (van Vuuren et al., 2011). The dependence of metrics on background conditions also
379 implies that metrics values inevitably change over time as background concentrations change as a
380 result of human activities, posing a challenge for the consistency of LCIA results with emissions
381 that take place over time. For instance, Reisinger et al. (2011) have found a 100-year GWP value
382 20% higher than today for methane under the lowest RCP for an emission occurring in 2100 and
383 10% lower by mid-century than today under the highest RCP. This further exemplifies that LCA
384 practitioners cannot avoid subjective judgments when choosing a metric; they can only assess the
385 consequences of alternative choices and communicate to end-users whether the results of LCA
386 are robust across a wide range of different metric approaches or highly contingent on particular
387 choices.

388 **3.3. Climate-carbon cycle feedbacks**

389 In the IPCC 4th AR, climate-carbon cycle feedbacks were included in the calculation of AGWP
390 for CO₂ but not for other GHGs (Forster et al., 2007). This inconsistency led to an
391 underestimation of GWP and even more for GTP values for non-CO₂ GHGs relative to CO₂,
392 because the warming caused by emission of a non-CO₂ gas causes CO₂ already in the atmosphere
393 at this time to persist for longer and thus add to the total warming effect caused by the non-CO₂
394 emission (Myhre et al., 2013). Indeed, the consideration of these feedbacks may have a
395 significant impact on emission metrics values. For instance, Gillet and Matthews (2010) found an
396 increase of 20% in GWP for CH₄ and N₂O (results were similar for both gases) and 80% in GTP
397 for CH₄ for a 100-year time horizon when adding climate-carbon cycle feedbacks. Collins et al.
398 (2013) found that the climate-carbon cycle feedbacks approximately double methane GTP for 100
399 years. In its 5th AR, the IPCC provides tentative values for illustration with and without

400 considering climate-carbon cycle feedbacks for some non-CO₂ forcers, with the 100-year GWP
401 increasing by 12 and 21% for N₂O and CH₄, respectively, and the 100-year GTP increasing by
402 27% for N₂O and more than doubling for CH₄, if climate-carbon cycle feedbacks are included.
403 Furthermore, there are also other feedbacks that can, for some forcers, be significant and that are
404 often not included in the calculation of GWPs. For example, these feedbacks relate to
405 atmospheric chemistry interactions (Shindell et al., 2009) and tropospheric O₃-carbon cycle
406 interactions (Collins et al., 2010).

407 Feedback mechanisms are very complex and can increase the uncertainty in a metric value. For
408 instance, the uncertainty in AGWP associated with climate-carbon cycle feedbacks may reach
409 ±100% of its best estimate (Myhre et al., 2013). In other words, uncertainties related to feedback
410 effect are comparable in magnitude to the strength of the feedback itself. The consideration of
411 these feedbacks thus results in a trade-off between accuracy, consistency and comprehensiveness,
412 but there is a clear case that climate-carbon cycle feedbacks should be treated consistently as they
413 are a well-understood (even if difficult to quantify) feature of the climate system.

414 **3.4. Regional variations**

415 The global mean temperature change depends on the location of emissions for NTCFs. These
416 regional variations may be addressed using different metric values for region of emission (e.g.
417 Berntsen et al., 2005; Stohl et al., 2015). Indeed, climate impacts of WMGHGs do not depend on
418 the location of emissions because they have lifetimes long enough so that they get well mixed in
419 the troposphere. However, NTCFs have much shorter lifetimes and impacts depend on where
420 emissions occur. Metrics for NTCFs may thus be given for region of emission. The IPCC 5th AR
421 presents the results of GWP and GTP for NO_x, CO, VOC, black carbon and organic carbon from
422 different studies. For instance, Fry et al. (2012) and Collins et al. (2013) have calculated GWP
423 and GTP values for 20- and 100-year time horizons for four regions (East Asia, European Union


424 and North Africa, North America, and South Asia) for NO_x, CO and VOC. For NO_x, GWP varies
425 from -40.7 to 6.4 kgCO₂-eq/kg for 20 years and from -25.3 to -5.3 kgCO₂-eq/kg for 100 years,
426 showing the influence of the location of emission on the results. The results obtained by Shindell
427 & Faluvegi (2009), Fuglestvedt et al. (2010), Bond et al. (2011, 2013), and Aamaas et al. (2015,
428 2016) also show a high regional variability for GWP and GTP of NTCFs.

429 On the other hand, the climate response also varies from one region to another. These regional
430 variations may be addressed using different metric values for region where climate impacts occur
431 (Collins et al., 2013; Shindell & Faluvegi, 2009). Some researchers have worked on the
432 development of metrics that take into account the regional variability of climate impacts (e.g.
433 Collins et al., 2013; Lund et al., 2012; Shindell, 2012) and the application of these metrics has
434 begun (Lund et al., 2014; Stohl et al., 2015; Sand et al., 2016). The conclusions from the IPCC 5th
435 AR are that additional studies are still needed to ensure their robustness (Myhre et al., 2013).

436 **3.5. The time dimension**

437 The time horizon defines the length of time over which impacts of climate forcers are integrated
438 for cumulative metrics, or the number of years into the future at which an instantaneous metric is
439 evaluated. Fixed time horizons are usually applied in LCA. This means that impacts are assessed
440 over a fixed period of time (e.g. 100 years) following each emission. The use of a fixed time
441 horizon for GWP ensures equal inclusion and weighting of actual impacts from emissions,
442 regardless of when in a product life cycle they occur (Peters et al., 2011b; Jørgensen &
443 Hauschild, 2013). Impacts are thus assessed over a sliding time window if emissions are spread
444 over several years as shown in Figure 3a. On the other hand, the use of a variable time horizon
445 depending on the relative timing of life cycle emissions allows the assessment of climate change
446 impacts with regard to a fixed future reference time as shown in Figure 4b. This approach implies
447 that only impacts up to the fixed point in time are deemed relevant, which may be relevant in

448 some decision contexts. For instance, it has been used in recent literature in the dynamic LCA
 449 approach and for the calculation of payback times for land use change emissions mitigation
 450 (Levasseur et al., 2010; O’Hare et al., 2009) or for the computation of emission metrics for
 451 biogenic emissions from long rotation biomass amongst others (Cherubini et al., 2016; Guest et
 452 al., 2013; Schivley et al., 2015).



453

454 Figure 3. Fixed time horizons (Sliding time window) (a) are usually applied in LCA. Variable
 455 time horizons (Fixed end point in time) (b) may also be relevant in some decision contexts

456 3.5.1. Choice of a time horizon

457 LCA practitioners can choose different time horizons, leading to different metric values. For
 458 instance, the value of GWP for short-lived WMGHGs and NTCFs decreases with increasing time
 459 horizon as the integrated radiative forcing of the reference gas CO_2 in the denominator continues
 460 to increase, as shown in Figure 8.29 of the IPCC 5th AR (Myhre et al., 2013). The 100-year time
 461 horizon is applied most often because of its adoption for the Kyoto Protocol. However, Shine
 462 (2009) argues that a 100-year time horizon was selected as an “inadvertent consensus” probably
 463 because it was the middle value between those used by the IPCC for its calculations (20, 100, and
 464 500 years). The use of this time horizon is thus not scientifically more justified than any other
 465 time horizon, but rather a value-based choice that must be based on considerations that include
 466 both science and ethics (Fuglestvedt et al., 2003; Levasseur et al., 2012; Shine, 2009; Tanaka et

467 al., 2010). For instance, the use of a 500-year time horizon has the advantage of indicating the
468 persistence and long-term warming effect of some gases, and selecting a shorter time horizon
469 may violate the principle of inter-generational equity widespread in LCA, because future impacts
470 beyond the time horizon are ignored entirely (Brandão et al., 2013).


471 On the other hand, a longer time horizon relies on the modelling of atmospheric or climate
472 processes that will occur far in the future, leading to higher uncertainties, and the relevance of a
473 midpoint metric for a future society several hundred years into the future is very difficult to
474 quantify. In its 5th AR, the IPCC does not provide values for a 500-year time horizon due to large
475 uncertainties and strong assumptions of constant background conditions, as well as ambiguity
476 related to what the metrics indicate on such timescales, which is especially the case for GWP
477 (Myrhe et al., 2013).

478 As shown in Figure 4, the selection of a time horizon strongly affects the value of metrics,
479 especially for climate forcers with a lifetime of less than roughly 100 years (Smith et al., 2012). It
480 is thus critically important to understand the underlying implications of the choice of time
481 horizon. Figure 4 shows how the instantaneous forcing, cumulative forcing, and temperature
482 change profiles of CO₂ and CH₄ pulse emissions evolve over time (for the sake of presentation,
483 CO₂ emissions are adjusted to match CH₄ emissions in magnitudes in terms of GWP100). At a
484 time horizon of 40 years, CO₂ has a higher instantaneous forcing and lower cumulative forcing,
485 and the temperature change of the two emissions is nearly identical (given an emissions pulse of
486 CO₂ 28 times larger than that of CH₄). This is, of course, different from the conclusion that
487 would be reached with a time horizon of 20 or 100 years. The wide range in metric values for a
488 relatively short-lived WMGHG such as methane reflects a fundamental difference in how
489 different components affect the climate over time. Methane has a strong short-term forcing that
490 almost disappears after a few decades as it leaves the atmosphere. In contrast, CO₂ has a more
491 persistent effect, leading to greater impacts than methane in the more distant future.

492 A short time horizon will put emphasis on short-term impacts (or rate of change) and thus gives a
493 higher relative importance to short-lived WMGHGs and NTCFs, while a longer time horizon will
494 do the opposite. The choice of a time horizon for instantaneous metrics can be aligned with, in the
495 case of GTP, the estimated peaking year of the global mean surface temperature. In that case, this
496 would reflect a primary goal of limiting peak warming. However, for LCAs that consider
497 emissions occurring over several years, fixing the end of the time horizons to a given calendar
498 year requires the use of a variable time horizon for climate metrics depending on the timing of
499 each emission relative to that calendar year as shown in Figure 3b.

500 **3.5.2. Discount rates**

501 Some methods have also been proposed to develop emission metrics while avoiding the use of a
502 time horizon entirely (e.g. Boucher, 2012; Cherubini et al., 2014; Smith et al., 2012; Wigley,
503 1998). Some forms of discounting are usually present in such metrics and the selection of any
504 discount rate requires value judgments and cannot be based on science alone, as the economic
505 literature amply attests (e.g. Anthoff et al., 2009; Goulder & Williams, 2012). This choice of
506 discount rate is in terms of value judgements similar to the choice of time horizon when using
507 cumulative metrics. The selection of a fixed time horizon for cumulative metrics is thus a
508 particular case of discounting using a 0% rate for impacts occurring prior to the end of the time
509 period, and an infinite discount rate for impacts occurring beyond, which are still significant for
510 long-lived gases such as CO₂ (IPCC, 2014a; Myhre et al., 2013; Tol et al., 2012). At the same
511 time, selection of an instantaneous metric such as GTP evaluates warming only in one specific
512 year and thus discounts entirely the warming for any other year before or after this target year
513 (Tol et al., 2012).



514

515 Figure 4. Instantaneous radiative forcing, cumulative radiative forcing, and instantaneous
 516 temperature change over time caused by emissions of 28 kg CO_2 and 1 kg non-fossil CH_4

517 **3.5.3. Considering the timing of emissions**

518 Approaches have been proposed to compare LCA results using a fixed end point in time for
 519 global warming impacts. All of them are based on the use of time horizon-dependent CFs to take
 520 into account the number of years occurring between the emission to assess and the chosen end-
 521 time for the analysis following the approach shown in Figure 3b (Cherubini et al., 2011; Kendall,
 522 2012; Levasseur et al., 2010). The use of such methods implies the elaboration of temporal
 523 emission profiles compared to aggregated emissions, as currently done in LCA, because one
 524 needs to know the time elapsed between the each emission and the fixed end point in time
 525 selected to use the right characterization factors. Moreover, as explained by Jørgensen and
 526 Hauschild (2013), using a fixed end point in time as shown in Figure 3b makes emissions
 527 occurring later have less impact, compared to similar emissions occurring earlier. However, these
 528 approaches are useful to look at the evolution of impacts through time such as done with dynamic
 529 LCA (Levasseur et al., 2010) or when only impacts until a specific point in time are deemed
 530 relevant.

531 Timing is also critical when assessing the climate impacts of emissions of different forcers in
 532 relation to a climate stabilisation limit, e.g. the $2^\circ C$ limit. The GWP has been criticised as being
 533 economically inefficient and indeed inconsistent with the ultimate objective of the UNFCCC to

534 stabilise greenhouse gas concentrations at a given level (Manne & Richels, 2001; Myhre et al.,
535 2013; Shine et al., 2007; Johansson, 2012; Tol et al., 2012). When assessing emissions in relation
536 to its contribution to remain within a pre-defined limit, it is therefore essential to have an estimate
537 of the year in which the temperature threshold will be reached. A time-dependent GTP was
538 therefore proposed to deal with this situation (Shine et al., 2007), where the metric time horizon is
539 based on the time left to the year the temperature peak is reached (Manne & Richels, 2001;
540 Michaelis, 1992; Shine et al., 2007). Other approaches, where the metric time horizon is based on
541 the time left to the year a prescribed limit is reached, exist as well, e.g. the climate tipping
542 potential (CTP) proposed by Jørgensen et al. (2014). For this approach, a ‘planetary boundary’
543 context is included, as the absolute, cumulative impacts of GHG emissions, up until the year in
544 which a predefined limit will be reached, are expressed as a fraction of the ‘capacity’ left before
545 exceeding the predefined limit. However, as shown by Persson et al. (2015), adopting a time-
546 dependent metric implies a commitment to significantly higher metric values for NTCFs and
547 short-lived WMGHGs as we get closer to the time when peak temperatures are reached. To avoid
548 disregarding longer term impacts beyond the chosen time horizon, an approach that both consider
549 the time until a threshold is reached and the climate impacts beyond that point in time might be
550 useful (Johansson, 2012; Jørgensen et al., 2014; Smith et al., 2012).

551 **4. Recommendations**

552 The use of GWP with a fixed time horizon has come under increased scrutiny as awareness of its
553 limitations has become more widespread over the recent past. GWP was the only metric
554 presented and discussed in the IPCC First AR. The IPCC 4th AR was the first to introduce and
555 discuss an alternative metric, i.e., GTP, but still considered that GWP was a “useful metric for
556 comparing the potential climate impact of the emissions of different [long-lived gases]“ (Forster
557 et al., 2007). The IPCC 5th AR shows an evolution in the thinking, clearly stating that the
558 selection of a metric depends on the policy goal to achieve, and that different metrics may lead to

559 different valid conclusions about the relative importance of emissions (Myhre et al., 2013). For
560 instance, Allen et al. (2016) showed that the GWP100 measures the relative impact of both
561 cumulative species and short-lived climate forcers on realized warming 20-40 after the time of
562 emissions. This metric is thus less suitable to account for long-term climate change impacts. In
563 this section, we first provide recommendations for a more robust assessment and interpretation of
564 potential climate change impacts in LCA following the most recent climate research findings
565 presented in this review. However, since current LCIA methods use cumulative metrics such as
566 GWP and do not account for the timing of emissions, we then present some recommendations
567 regarding the integration of the most recent findings of the IPCC 5th AR and concurrent scientific
568 research in LCA practice.

569 **4.1. Improving climate change impact assessment in LCA**

570 Emission metrics are used to help decision makers identify how emissions of different climate
571 forcers compare in terms of impacting specific aspects of climate change. The appropriateness of
572 a climate change metric for a given application thus depends on the purpose that it is meant to
573 serve i.e., the overall goal of climate policies and which aspects of climate change are deemed
574 relevant (Fuglestvedt et al., 2003; Myhre et al., 2013). In an LCA context, we consider it critical
575 that practitioners assess whether their results depend strongly on the choice of metric. This will
576 depend on the purpose an LCA is intended to serve (i.e. comparison of carbon footprints of two
577 products, or long-term company-wide mitigation strategy). The choice of metric thus cannot be
578 made independently of the values, goals, and scope of the end-user of an LCA.

579 In practice, it can make sense to use several complementary metrics that will serve different
580 purposes, and from there allow LCA practitioners to get a better understanding about the
581 robustness of the LCA study to different metrics during the interpretation phase of LCA. LCA
582 methodology and software should gradually move towards a situation where the state-of-the-art is

583 represented by the use of several sets of CFs based on different metrics and/or time horizons.
584 Analysts can then test how robust the overall conclusions are with respect to such different
585 choices. If results differ significantly for different metrics, they can argue or demonstrate why one
586 metric choice would be preferable to the others for a given purpose. The range of results from
587 different metrics should become part of communicating the ambiguity and uncertainty of LCA
588 results (where ‘uncertainty’ reflects the scientific uncertainty and ‘ambiguity’ the dependence on
589 human choices in methodologies or purposes of the LCA). This will certainly require an initial
590 transition phase for adaptation of practitioners and updates of CFs in the common databases and
591 software providers for LCA analyses. However, the availability of characterization factors in the
592 IPCC 5th AR makes this transition easier, and relatively little adaptation efforts will ensure an
593 important step forward in the robustness of climate change impact assessment in LCA. We
594 believe that this increased transparency about the hidden value judgements in LCA is critical to
595 ensure that an LCA actually serves the intended purpose.

596 The use of multiple indicators for assessing climate change impacts in other branches derived
597 from LCA like carbon footprints and product labels can be more challenging. The different
598 groups and stakeholders should start a debate on how to reflect these considerations in the
599 different applications. The alternative, if they simply continue to rely on a ‘preferred’ metric
600 independently of its meaning and end user goals, this could lead to perverse outcomes. For
601 example, imagine a consumer keen to support the rapid reduction of CO₂ emissions, consistent
602 with the finding by the IPCC that CO₂ emissions have to drop to zero before 2100 to limit
603 warming to 2°C. Faced with the choice of whether to purchase product A or product B, the
604 consumer will rely on the reported carbon footprint. But product A might have a very large CH₄
605 component while product B may release almost exclusively CO₂. Product A may have a larger
606 carbon footprint in the metric that practitioners have decided to use, and the consumer would
607 therefore purchase product B, even though this results in greater CO₂ emissions that lead to long-

608 term climate change. The extent to which the complexity of alternative choices can be
609 communicated successfully to end-users will of course depend strongly on the context; while it is
610 unlikely to be successful for product labels in supermarkets, such a conversation should be
611 entirely feasible and in fact an obligation where LCA practitioners interact with individual clients
612 from industry or government (including, in fact, the owners of supermarket chains that decide on
613 what labelling system they wish to use and communicate).

614 Based on these findings, we recommend the following step-by-step approach:

- 615 a) Selection of a few metrics and time horizons that differ in the type of climate response
616 they capture for both WMGHGs and NTCFs. In addition to using GWP100 for
617 comparability purposes, and potentially one or several other well thought-through choices
618 of metrics and time horizons based on the goal and scope of the study, we recommend to
619 perform a sensitivity analysis. For this purpose, GWP20 and GTP100 may provide a
620 suitable range of metric values as they allow LCA users to explore their results using
621 either a short-term perspective with very high (GWP20) or a long-term perspective with
622 very low (GTP100) weighting for short-lived WMGHGs and NTCFs.
- 623 b) Calculation of impact scores using the selected metrics to get a sense of whether the
624 results are critically dependent on the metric choice.
- 625 c) If the conclusions of the LCA study (relative to the specific purpose it seeks to serve) do
626 not change significantly using different metrics or time horizons, there is no need to
627 communicate on several metrics.
- 628 d) If the conclusions of the LCA study vary using different metrics or time horizons, the
629 results must be communicated very carefully, explaining how the answer differs between
630 metrics and time horizons and why, and guiding users as to whether a particular metric
631 choice may be more suitable than others for the particular purpose and ultimate goals that
632 the end-user may have in mind.

633 Some choices must be made when selecting emission metrics regarding the characteristics and
634 modelling conditions presented in the previous sections: type of effect modelled (radiative
635 forcing, temperature, etc.), instantaneous vs. cumulative, time horizon and/or discounting,
636 constant vs. variable background, etc. Some of these choices are scientific, while others are more
637 policy-related and cannot be based solely on scientific studies (Tanaka et al., 2010). The choice of
638 metric type and time horizon as proposed in this step-by-step approach may have much larger
639 effects on decisions than the improvement of input parameters to the metrics (Myhre et al., 2013).

640 **4.2. Updating GWP and GTP according to IPCC 5th AR**

641 Working Group I produced a long list of updated GWP and GTP values for 20- and 100-year time
642 horizons and GTP values for 50-year time horizon in the IPCC 5thAR for more than 200 GHGs,
643 including the quantification of uncertainties. We recommend that these values (or updated values
644 from a more recent IPCC AR when available) are used in LCA if GWP or GTP are selected as
645 climate change midpoint indicator. The following recommendations regard the consideration of
646 climate-carbon cycle feedbacks for non-CO₂ GHGs, and the development of CFs for NTCFs.

647 **4.2.1. Consideration of climate-carbon cycle feedbacks for non-CO₂ components**

648 As explained in section 3.3, the use of GWP values including climate-carbon cycle feedbacks for
649 CO₂ but not for other GHGs is inconsistent and leads to lower relative impacts for non-CO₂
650 GHGs. However, high uncertainties are associated with the inclusion of feedbacks for non-CO₂
651 GHGs because there are still just a few values available in the literature. Research is still needed
652 to improve the reliability of these GWP values. We thus recommend including climate-carbon
653 cycle feedbacks for all GHGs, providing associated uncertainty values with CFs, and performing
654 an uncertainty analysis. This could be done by simple estimates based on the uncertainty ranges
655 for metric values or using a Monte Carlo method. If this cannot be done, we recommend
656 calculating impact scores with and without the inclusion of climate-carbon cycle feedbacks for

657 non-CO₂ GHGs to determine if their inclusion changes conclusions significantly. If it is the case,
658 we recommend discussing the importance of this choice when communicating results (this
659 process is similar to steps b-d in the above recommended steps for evaluation of different
660 metrics).

661 As a rule of thumb, the inclusion of climate-carbon cycle feedbacks tends to have a smaller effect
662 on metric values for NTCFs and short-lived WMGHGs than alternative metric choices (e.g. GTP
663 instead of GWP) and time horizons. If the sensitivity test regarding alternative metrics shows
664 little effect on the overall LCA result, then testing for the sensitivity of climate-carbon cycle
665 feedbacks could perhaps be omitted.

666 **4.2.2. Development of CFs for NTCFs**

667 As discussed in section 3.4, climate impacts of NTCFs depend on the location of emissions.
668 Global scale metrics may thus not be the most appropriate. Moreover, NTCFs are tightly coupled
669 to the hydrological cycle and atmospheric chemistry which are very complex processes difficult
670 to model and validate. The values associated to NTCF metrics are more uncertain than for
671 WMGHGs and there are substantial variations across studies (Myhre et al., 2013).

672 The quality of the metrics of NTCFs may improve over the next years as research goes on, even
673 though they will always be more uncertain and sensitive to assumptions than the metrics of
674 WMGHGs. However, for some sectors or activities, NTCFs can make an important contribution
675 to the climate impact category (e.g. Fuglestvedt et al, 2008). We thus recommend calculating
676 impact scores with and without NTCFs using global average values (i.e. not specific to selected
677 regions) from the literature as presented in the IPCC 5th AR. Once again, if their consideration
678 changes conclusions, we recommend discussing it when communicating results, explaining
679 uncertainty and regional variability issues. If the location of emissions is known and a
680 regionalized life cycle assessment approach is used, we recommend using region-specific GWP

681 and GTP values for NTCFs to reduce uncertainties. Indeed, if the location of emissions is known,
682 it is conceptually easy to use emission-region-specific CFs that express the global climate impact
683 of NTCFs, although not all LCA software or life cycle inventory databases support this feature
684 yet.

685 Some NTCFs (e.g. SO₂ and organic carbon) have negative metric values because of their cooling
686 effect when in the atmosphere as illustrated in FAQ 8.2 Figure 1 of the IPCC 5th AR (Myhre et
687 al., 2013). Globally, the combined cooling effect is significant and considered to have offset some
688 of the warming from WMGHGs that would otherwise have occurred. However, concerns may be
689 raised regarding the inclusion of pollutants with a cooling effect in the global warming impact
690 category because they may cause other type of environmental impacts such as acidification for
691 the case of SO₂. This means that one could favour a high GHG and high SO₂ emitter compared to
692 a lower GHG and no SO₂ emitter if only the global warming LCA result is considered, leading to
693 higher other environmental impacts such as acidification or human health issues. This example
694 shows why it is crucial to use multiple indicators, as LCA usually does, to guide choices. We thus
695 recommend including climate forcers with negative metric values if NTCFs are considered at all
696 for consistency purposes. However, the warming and cooling effects should be presented
697 separately to facilitate transparency and analysis of results.

698 In conclusion, given the many relevant metrics and the broad set of emission components and
699 effects acting on very different timescales, LCA studies may benefit from moving away from
700 single-metric studies towards a multi-metric perspective and sensitivity tests, combined with (or
701 followed by) a careful exchange with the end-users of LCA to ensure that those value judgements
702 that have to be made serve the intended purposes of those who use the resulting information. This
703 will help to communicate the complexity of the system and processes and at the same time
704 increase the transparency regarding value related choices inherent to LCA results.

705 **Acknowledgments.**

706 This work has been done on a voluntary basis through the Global Warming Task Force of the
707 project ‘Global Guidance on Environmental Life Cycle Assessment Indicators’ of the United
708 Nations Environment Programme (UNEP) / Society of Environmental Toxicology and Chemistry
709 (SETAC) Life Cycle Initiative. Public and private sector sponsors are listed on the Initiative’s
710 website (www.lifecycleinitiative.org). The views expressed in this article are those of the authors
711 and do not necessarily reflect those of the various affiliated organizations. DJAJ contribution was
712 financed by the Swedish Energy Agency.

713 **Note.**

714 The first author of this article is the chair of the task force and main author of this paper. The last
715 author is the co-chair of the task force. Other authors are members of the task force and are
716 ordered alphabetically.

717 **References.**

718 Aaheim, A.; Fuglestvedt, J.S.; Godal, O. Costs savings of a flexible multi-gas climate policy.
719 *Energ. J.* **2006**, *27*, 485-501.

720 Aamaas, B.; Berntsen, T.K.; Fuglestvedt, J.S.; Shine, K.P.; Bellouin, N. Multimodel emission
721 metrics for regional emissions of short lived climate forcers. *Atmos. Chem. Phys.* **2015**, *15*,
722 26089-26130.

723 Aamaas, B.; Berntsen, T.K.; Fuglestvedt, J.S.; Shine, K.P., Bellouin, N. Regional emission
724 metrics for short-lived climate forcers from multiple models. *Atmos. Chem. Phys.* **2016**, *16*, 7451-
725 7468.

726 Allen, M.R. Fuglestedt, J.S., Shine, K.P, Reisinger, A., Pierrehumbert, R.T., Forster, P.M. New
727 use of global warming potentials to compare cumulative and short-lived climate pollutants. *Nat.*
728 *Clim. Change.* **2016**, .

729 Anthoff, D.; Tol, R.S.J.; Yohe, G.W. Discounting for climate change. *Economics.* **2009**, *3* (2009-
730 24), 1-22.

731 Azar, D.; Johansson, D.J.A. On the relationship between metrics to compare greenhouse gases –
732 the case of IGTP, GWP and SGTP. *Earth Syst. Dynam.* **2012**, *3* (2), 113-141.

733 Baker, A.C.; Glynn, P.W.; Riegl, B. Climate change and coral reef bleaching: an ecological
734 assessment of long-term impacts, recovery trends and future outlook. *Estuar. Coast. Shelf S.*
735 **2008**, *80* (4), 435-471.

736 Berntsen, T.K.; et al. Climate response to regional emissions of ozone precursors: sensitivities
737 and warming potentials. *Tellus B.* **2005**, *57* (4), 283-304.

738 Bond, T.; Zarzycki, C.; Flanner, M.; Koch, D. Quantifying immediate radiative forcing by black
739 carbon and organic matter with the Specific Forcing Pulse. *Atmos. Chem. Phys.* **2011**, *11* (4),
740 1505-1525.

741 Bond, T.; et al. Bounding the role of black carbon in the climate system: A scientific assessment.
742 *J. Geophys. Res.-Atmos.* **2013**, *118* (11), 5380-5552.

743 Boucher, O.; Reddy, M. Climate trade-off between black carbon and carbon dioxide emissions.
744 *Energ. Policy.* **2008**, *36* (1), 193-200.

745 Boucher, O. Comparison of physically- and economically-based CO₂-equivalences for methane.
746 *Earth Syst. Dynam.* **2012**, *3* (1), 49-61.

747 Bowerman, N.H.A.; Frame, D.J.; Huntingford, C.; Lowe, J.A.; Smith, S.M.; Allen, M.R. The role
748 of short-lived climate pollutants in meeting temperature goals. *Nat. Clim. Change*. **2013**, *3*, 1021-
749 1024.

750 Brandão, M; et al. Key issues and options in accounting for carbon sequestration and temporary
751 storage in life cycle assessment and carbon footprinting. *Int. J. Life Cycle Ass.* **2013**, *18* (1), 230-
752 240.

753 Cherubini, F.; Peters, G.P.; Berntsen, T.; Strømman, A.H.; Hertwich, E. CO₂ emissions from
754 biomass combustion for bioenergy: atmospheric decay and contribution to global warming. *GCB*
755 *Bioenergy*, **2011**, *35*, 413-426.

756 Cherubini, F.; Bright, R.; Strømman, A.H. Global climate impacts of forest bioenergy: what,
757 when and how to measure? *Environ. Res. Lett.* **2013**, *8*, 014049.

758 Cherubini, F.; Gasser, T.; Bright, R.; Ciais, P.; Strømman, A.H. Linearity between temperature
759 peak and bioenergy CO₂ emission rates. *Nat. Clim. Change*. **2014**, *4*, 983-987.

760 Cherubini, F., Huijbregts, M., Kindermann, G., Van Zelm, R., Van Der Velde, M., Stadler, K.,
761 Strømman, A.H. Global spatially explicit CO₂ emission metrics for forest bioenergy. *Scientific*
762 *Reports*, **2016**, *6*, 20186.

763 Collins, W.J.; Sitch, S.; Boucher, O. How vegetation impacts affect climate metrics for ozone
764 precursors. *J. Geophys. Res.* **2010**, *115* (D23), D23308.

765 Collins, W.J.; Fry, M.M.; Yu, H.; Fuglestedt, J.S.; Shindell, D.T.; West, J.J. Global and regional
766 temperature-change potentials for near-term climate forcers. *Atmos. Chem. Phys.* **2013**, *13* (5),
767 2471-2485.

768 Deuber, O.; Luderer, G.; Edenhofer, O. Physico-economic evaluation of climate metrics: A
769 conceptual framework. *Environ. Sci. Policy*. **2013**, *29*, 37-45.

770 Eckaus, R.S. Comparing the effects of greenhouse gas emissions on global warming. *Energy J.*
771 **1992**, *13* (1), 25-36.

772 Edwards, M.R.; Trancik, J.E. Climate impacts of energy technologies depend on emissions
773 timing. *Nat. Clim. Change*. **2014**, *4*, 347-352.

774 Ekholm, T.; Lindroos, T.J.; Savolainen, I. Robustness of climate metrics under climate policy
775 ambiguity. *Environ. Sci. Policy*. **2013**, *31*, 44-52.

776 Forster, P.; et al. Changes in Atmospheric Constituents and in Radiative Forcing. In *Climate*
777 *Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth*
778 *Assessment Report of the Intergovernmental Panel on Climate Change*; Solomon, S., et al. Eds.;
779 Cambridge University Press: Cambridge and New-York, NY, **2007**; pp 129-234.

780 Friedlingstein, P.; et al. Climate-carbon cycle feedback analysis: Results from the C⁴MIP model
781 intercomparison. *J. Climate*. **2006**, *19* (14), 3337-3353.

782 Fry, M. et al. The influence of ozone precursor emissions from four world regions on
783 tropospheric composition and radiative climate forcing. *J. Geophys. Res.-Atmos.* **2012**, *117* (D7),
784 D07306.

785 Fuglestedt, J.; Berntsen, T.; Godal, O.; Sausen, R.; Shine, K.; Skodvin, T. Metrics of climate
786 change: Assessing radiative forcing and emission indices. *Clim. Change*. **2003**, *58* (3), 267-331.

787 Fuglestedt, J.; Berntsen, T.; Myhre, G.; Rypdal, K.; Skeie, R.B. Climate forcing from the
788 transport sectors. *P. Natl. Acad. Sci. USA*. **2008**, *105* (2), 454-458.

789 Fuglestad, J.S.; et al. Transport impacts on atmosphere and climate: Metrics. *Atmos. Environ.*
790 **2010**, *44* (37), 4648-4677.

791 Geoffroy, O.; Saint-Martin, D.; Oliv  , D.J.L.; Voldoire, A.; Bellon, G.; Tyt  ca, S. Transient
792 climate response in a two-layer energy-balance model. Part I: Analytical solution and parameter
793 calibration using CMIP5 AOGCM experiments. *J. Clim.* **2013**, *26* (6), 1841-1857.

794 Gillet, N.P.; Matthews, H.D. Accounting for carbon cycle feedbacks in a comparison of the
795 global warming effects of greenhouse gases. *Environ. Res. Lett.* **2010**, *5*, 034011.

796 Goedkoop, M.; Heijungs, R.; Huijbregts, M.; De Schryver, A.; Struijs, J.; van Zelm, R. *ReCiPe*
797 *2008. A life cycle impact assessment method which comprises harmonised category indicators at*
798 *the midpoint and endpoint level*; First edition (version 1.08), **2013**. [http://www.lcia-](http://www.lcia-recipe.net/file-cabinet)
799 [recipe.net/file-cabinet](http://www.lcia-recipe.net/file-cabinet).

800 Goulder, L.H.; Williams, R.C.III. The choice of discount rate for climate change policy
801 evaluation. *Clim. Change Econ.* **2012**, *3* (4), 1250024.

802 Guest, G., Cherubini, F., & Str  mman, A.H. The role of forest residues in the accounting for the
803 global warming potential of bioenergy. *GCB Bioenergy*, **2013**, *5*, 459-466.

804 Hammit, J.K.; Jain, A.K.; Adams, J.L.; Wuebbles, D.J. A welfare-based index for assessing
805 environmental effects of greenhouse-gas emissions. *Nature*. **1996**, *381*, 301-303.

806 Herrington, T.; Zickfeld, K. Path independence of climate and carbon cycle response over a broad
807 range of cumulative carbon emissions. *Earth Syst. Dynam.* **2014**, *5* (2), 409-422.

808 Huang, C.R.; Barnett, A.G.; Wang, X.M.; Vaneckova, P.; Fitzgerald, G.; Tong, S.L. Projecting
809 future heat-related mortality under climate change scenarios: a systematic review. *Environ.*
810 *Health Persp.* **2011**, *119* (12), 1681-1690.

811 IPCC. *Climate Change: The Intergovernmental Panel on Climate Change Scientific Assessment*;
812 Cambridge University Press: Cambridge, **1990**.

813 IPCC. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to*
814 *the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; Pachauri, R.K.,
815 Meyer, L.A. Eds.; IPCC: Geneva, **2014a**.

816 IPCC. *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and*
817 *Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the*
818 *Intergovernmental Panel on Climate Change*; Field, C.B., et al. Eds.; Cambridge University
819 Press: Cambridge and New-York, NY, **2014b**.

820 *ISO 14044 - Environmental management – Life cycle assessment – Requirements and guidelines*;
821 International Organization for Standardization: Lausanne, **2006**.

822 Johansson, D.J.A.; Persson, U.M.; Azar, C. The cost of using global warming potentials:
823 Analysing the trade off between CO₂, CH₄ and N₂O. *Clim. Change*. **2006**, 77 (3-4), 291-309.

824 Johansson, D.J.A. Economics- and physical-based metrics for comparing greenhouse gases. *Clim.*
825 *Change*. **2012**, 110 (1), 123-141.

826 Joos, F.; et al. Carbon dioxide and climate impulse response functions for the computation of
827 greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.* **2013**, 13 (5), 2793-2825.

828 Jørgensen, S.V.; Hauschild, M.Z. Need for relevant timescales when crediting temporary carbon
829 storage. *Int. J. Life Cycle Ass.* **2013**, 18 (4), 745-754.

830 Jørgensen, S.V.; Hauschild, M.Z.; Nielsen, P.H. Assessment of the urgent impacts of greenhouse
831 gas emissions – the climate tipping potential (CTP). *Int. J. Life Cycle Ass.* **2014**, 19 (4), 919-930.

832 Kendall, A. Time-adjusted global warming potentials for LCA and carbon footprints. *Int. J. Life*
833 *Cycle Ass.* **2012**, 17 (8), 1042-1049.

834 Kirschbaum, M.U.F. Climate-change impact potentials as an alternative to global warming
835 potentials. *Environ. Res. Lett.* **2014**, 9, 034014.

836 Kolstad, C.; et al. Social, Economic and Ethical Concepts and Methods. In *Climate Change 2014:*
837 *Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report*
838 *of the Intergovernmental Panel on Climate Change*; Edenhofer, O., et al. Eds.; Cambridge
839 University Press: Cambridge and New-York, NY, **2014**; pp 207-282.

840 Lashof, D.A., & Ahuja, D.R. Relative contributions of greenhouse gas emissions to global
841 warming. *Nature.* **1990**, 344(6266), 529-531.

842 Lauder, A.R.; et al. Offsetting methane emissions – An alternative to emission equivalence
843 metrics. *Int. J. of Greenh. Gas Con.* **2013**, 12, 419-429.

844 Lenton, T.M.; et al. Tipping elements in the Earth's climate system. *P. Natl. Acad. Sci. USA.*
845 **2008**, 105 (6), 1786-1793.

846 Levasseur, A. ; Lesage, P. ; Margni, M. ; Deschênes, L. ; Samson, R. Considering time in LCA:
847 dynamic LCA and its application to global warming impact assessment. *Environ. Sci. Technol.*
848 **2010**, 44 (8), 3169-3174.

849 Levasseur, A.; Brandão, M.; Lesage, P.; Margni, M.; Pennington, D.; Clift, R.; Samson, R.
850 Valuing temporary carbon storage. *Nat. Clim. Change.* **2012**, 2, 6-8.

851 Levasseur, A. Climate change. In *LCA Compendium – The Complete World of Life Cycle*
852 *Assessment – Life Cycle Impact Assessment*; Klöpffer, W., Curran M.A., Series Eds; Hauschild,
853 M., Huijbregts, M., Volume Eds.; Springer: Dordrecht, **2015**; pp 39-50.

854 Li, S.; Jarvis, A. Long run surface temperature dynamics of an A-OGCM: The HadCM3 4XCO₂
855 forcing experiment revisited. *Clim. Dynam.* **2009**, *33* (6), 817-825.

856 Lund, M.T.; Berntsen, T.; Fuglestvedt, J.S.; Ponater, M.; Shine, K.P. How much information is
857 lost by using global-mean climate metrics? An example using the transport sector. *Clim. Change.*
858 **2012**, *113* (3), 949-963.

859 Lund, M.T.; Berntsen, T.K.; Heyes, C.; Klimont, Z.; Samset, B.H. Global and regional climate
860 impacts of black carbon and co-emitted species from the on-road diesel sector. *Atmos. Environ.*
861 **2014**, *98*, 50-58.

862 Manne, A.S. Richels, R.G. An alternative approach to establishing trade-offs among greenhouse
863 gases. *Nature.* **2001**, *410*, 675-677.

864 Manning, M.; Reisinger, A. Broader perspectives for comparing different greenhouse gases.
865 *Philos. T. R. Soc. A.* **2011**, *369* (1943), 1891-1905.

866 Michaelis, P. Global warming: Efficient policies in the case of multiple pollutants. *Environ.*
867 *Resour. Econ.* **1992**, *2* (1), 61-77.

868 Myhre, G.; Fuglestvedt, J.S.; Berntsen, T.K.; Lund, M.T. Mitigation of short-lived heating
869 components may lead to unwanted long-term consequences. *Atmos. Environ.* **2011**, *45* (33),
870 6103-6106.

871 Myhre, G.D.; et al. Anthropogenic and Natural Radiative Forcing. In *Climate Change 2013: The*
872 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*
873 *Intergovernmental Panel on Climate Change*; Stocker, T.F., et al. Eds.; Cambridge University
874 Press: Cambridge and New-York, NY, **2013**; pp 659-740.

875 O'Hare, M.; Plevin, R.J.; Martin, J.I.; Jones, A.D.; Kendall, A.; Hopson, E. Proper accounting for
876 time increases crop-based biofuels' greenhouse gas deficit versus petroleum. *Environ. Res. Lett.*
877 **2009**, *42*, 024001.

878 Olivie D.J.L.; Peters, G.; Saint-Martin, D. Atmosphere response time scales estimated from
879 AOGCM experiments. *J. Clim.* **2012**, *25* (22), 7956-7972.

880 O'Neil, B.C. Economics, natural science, and the costs of global warming potentials. *Clim.*
881 *Change*. **2003**, *58* (3), 251-260.

882 Peck, S.C.; Teisberg, T.J. Optimal carbon emissions trajectories when damages depend on the
883 rate or level of global warming. *Clim. Change*. **1994**, *28* (3), 289-314.

884 Penner, J.E.; Prather, M.J.; Isaksen, I.S.A.; Flugestvedt, J.S.; Klimont, Z.; Stevenson, D.S. Short-
885 lived uncertainty? *Nat. Geosci.* **2010**, *3*, 587-588.

886 Persson, U.M.; Johansson, D.J.A.; Cederberg, C.; Hedenus, F.; Bryngelsson, D. Climate metrics
887 and the carbon footprint of livestock products: where's the beef? *Environ. Res. Lett.* **2015**, *10*,
888 034005.

889 Peters, G.P.; Aamaas, B.; Berntsen, T.; Fuglestvedt, J.S. The integrated global temperature
890 change potential (iGTP) and relationships between emission metrics. *Environ. Res. Lett.* **2011a**, *6*,
891 044021.

892 Peters, G.P.; Aamaas, B.; Lund, M.T.; Solli, C.; Fuglestvedt, J.S. Alternative "global warming"
893 metrics in life cycle assessment: A case study with existing transportation data. *Environ. Sci.*
894 *Technol.* **2011b**, *45* (20), 8633-8641.

895 Reisinger, A.; Meinshausen, M.; Manning, M.; Bodeker, G. Uncertainties of global warming
896 metrics: CO₂ and CH₄. *Geophys. Res. Lett.* **2010**, *37*, L14707.

897 Reisinger, A.; Meinshausen, M.; Manning, M. Future changes in global warming potentials under
898 representative concentration pathways. *Environ. Res. Lett.* **2011**, *6*, 024020.

899 Reisinger, A.; Havlik, P.; Riahi, K.; van Vliet, O.; Obersteiner, M.; Herrero, M. Implications of
900 alternative metrics for global mitigation costs and greenhouse gas emissions from agriculture.
901 *Clim. Change.* **2013**, *117* (4), 677-690.

902 Rilley, J.M.; Richards, K.R. Climate change damage and the trace gas index issue. *Environ.*
903 *Resour. Econ.* **1993**, *3* (1), 41-61.

904 Rogelj, J.; et al. Disentangling the effects of CO₂ and short-lived climate forcer mitigation. P.
905 *Natl. Acad. Sci. USA.* **2014**, *111* (46), 16325-16330.

906 Rotmans, J., & den Elzen, M.G.J. A model-based approach to the calculation of global warming
907 potentials (GWP). *Int. J. Climatol.*, **1992**, *12*, 865-874.

908 Sand, M., Berntsen, T.K., von Salzen, K., Flanner, M.G., Langner, J., Victor, D.G. Response of
909 Arctic temperature to changes in emissions to short-lived climate forcers. *Nat. Clim. Change.*
910 **2016**, *6*, 286-289.

911 Schivley, G.; Ingwersen, W.W.; Marriott, J.; Hawkins, T.R.; Skone, T.J. Identifying/quantifying
912 environmental trade-offs inherent in GHG reduction strategies for coal-fired power. *Environ. Sci.*
913 *Technol.* **2015**, *49* (13), 7562-7570.

914 Shindell, D.; Faluvegi, G. Climate response to regional radiative forcing during the twentieth
915 century. *Nat. Geosci.* **2009**, *2*, 294-300.

916 Shindell, D.T.; Faluvegi, G.; Koch, D.M.; Schmidt, G.A.; Unger, N.; Bauer, S.E. Improved
917 attribution of climate forcing to emissions. *Science.* **2009**, *326* (5953), 716-718.

918 Shindell, D.T. Evaluation of the absolute regional temperature potential. *Atmos. Chem. Phys.*
919 **2012**, *12* (17), 7955-7960.

920 Shindell, D.; et al. Simultaneously mitigating near-term climate change and improving human
921 health and food security. *Science*. **2012**, *335* (6065), 183-189.

922 Shine, K.P.; Fuglestedt, J.S.; Hailemariam, K.; Stuber, N. Alternatives to the global warming
923 potential for comparing climate impacts of emissions of greenhouse gases. *Clim. Change*. **2005**,
924 *68* (3), 281-302.

925 Shine, K.P.; Berntsen, T.K.; Fuglestedt, J.S.; Skeie, R.B.; Stuber, N. Comparing the climate
926 effect of emissions of short- and long-lived climate agents. *Philos. T. Roy. Soc. A*. **2007**, *365*
927 (1856), 1903-1914.

928 Shine K.P. The global warming potential—the need for an interdisciplinary retrieval. *Clim. Change*.
929 **2009**, *96* (4), 467–472.

930 Shine, K.P.; Allan, R.P.; Collins, W.J.; Fuglestedt, J.S. Metrics for linking emissions of gases
931 and aerosols to global precipitation changes. *Earth Syst. Dynam.* **2015**, *6*, 719-760.

932 Smith, S.M.; Lowe, J.A.; Bowerman, N.H.A.; Gohar, L.K.; Huntingford, C.; Allen, M.R.
933 Equivalence of greenhouse-gas emissions for peak temperature limits. *Nat. Clim. Change*. **2012**,
934 *2*, 535-538.

935 Smith, S.J.; Karas, J.; Edmonds, J.; Eom, J.; Mizrahi, A. Sensitivity of multi-gas climate policy to
936 emission metrics. *Clim. Change*. **2013**, *117* (4), 663-675.

937 Sterner, E.; Johansson, D.J.A.; Azar, C. Emission metrics and sea level rise. *Clim. Change*. **2014**,
938 *127* (2), 335-351.

939 Stohl, A.; et al. Evaluating the climate and air quality impacts of short-lived pollutants. *Atmos.*
940 *Chem. Phys.* **2015**, *15*, 10529-10566.

941 Strefler, J.; Luderer, G.; Aboumahboub, T.; Kriegler, E. Economic impacts of alternative
942 greenhouse gas emission metrics: a model-based assessment. *Clim. Change.* **2014**, *125* (3), 319-
943 331.

944 Tanaka, K.; O'Neil, B.C.; Rokityanskiy, D.; Obersteiner, M.; Tol, R.S.J. Evaluating global
945 warming potentials with historical temperature. *Clim. Change.* **2009**, *96* (4), 443-466.

946 Tanaka, K.; Peters, G.P.; Flugestvedt, J.S. Policy update: Multicomponent climate policy: why do
947 emission metrics matter? *Carbon Manage.* **2010**, *1* (2), 191-197.

948 Tanaka, K.; Johansson, D.J.A.; O'Neil, B.C.; Fuglestvedt, J.S. Emission metrics under the 2°C
949 climate stabilization target. *Clim. Change.* **2013**, *117* (4), 933-941.

950 Tol, R.S.J. Berntsen, T. K. O'Neil, B.C. Fuglestvedt, J.S. Shine, K.P. A unifying framework for
951 metrics for aggregating the climate effect of different emissions. *Environ. Res. Lett.* **2012**, *7*,
952 044006.

953 UNFCCC. *Workshop on common metrics to calculate the CO₂ equivalence of anthropogenic*
954 *greenhouse gas emissions by sources and removal by sinks,*
955 **2012**. http://unfccc.int/methods/other_methodological_issues/items/6737.php.

956 UNFCCC. *SBSTA-IPCC Special Event: Common metrics to calculate the carbon dioxide*
957 *equivalence of greenhouse gases,*
958 **2014**. http://unfccc.int/meetings/bonn_jun_2014/workshop/8245.php.

959 van den Berg, M.; Hof, A.F.; van Vliet, J.; van Vuuren, D.P. Impact of the choice of emission
960 metric on greenhouse gas abatement and costs. *Environ. Res. Lett.* **2015**, *10* (2), 024001.

- 961 van Vuuren, D.P.; et al. The representative concentration pathways: an overview. *Clim. Change*.
962 **2011**, *109* (1-2), 5-31.
- 963 Vermeer, M.; Rahmstorf, S. Global sea level linked to global temperature. *P. Natl. Acad. Sci.*
964 *USA*. **2009**, *106* (51), 21527-21532.
- 965 Webster, P.J.; Holland, G.J.; Curry, J.A.; Chang, H.R. Changes in tropical cyclone number,
966 duration, and intensity in a warming environment. *Science*. **2005**, *309* (5742), 1844-1846.
- 967 Wigley, T.M.L. The Kyoto Protocol: CO₂, CH₄, and climate implications. *Geophys. Res. Lett.*
968 **1998**, *25* (13), 2285-2288.