

1 1. Title Page

(a) Alterations in the metabolic and cardiorespiratory response to exercise in Huntington's Disease

(b) Authors and affiliations

J Steventon^{1,2,3}, *J. Collett*^{4,5}, *H. Furby*^{2,3}, *K. Hamana*⁶, *C. Foster*³, *P. O'Callaghan*⁷, *A. Dennis*⁸, *R. Armstrong*⁹, *A. H. Németh*⁹, *A. E. Rosser*¹⁰, *K. Murphy*^{1,3}, *L. Quinn*¹¹, *M. Busse*¹², *H. Dawes*^{4,13}

1. School of Physics and Astronomy, Cardiff University
2. NMHRI, School of Medicine, Cardiff University
3. CUBRIC, School of Psychology, Cardiff University
4. Centre for Movement, Occupation and Rehabilitation Sciences, OxINMAHR, Oxford Brookes University, Oxford, UK.
5. Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK.
6. School of Healthcare Sciences, Cardiff University
7. Cardiology Department, University Hospital of Wales, Cardiff
8. FMRIB Centre, Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, OX3 9DU, UK
9. Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, UK;
10. IPMCN, School of Medicine, Cardiff University
11. Department of Biobehavioral Sciences, Teachers College, Columbia University, USA.
12. Centre for Trials Research, Cardiff University, UK;
13. Department of Clinical Neurology, University of Oxford, Oxford, UK.

(c) **Corresponding author:** *Dr Jessica Steventon, Cardiff University Brain Research Imaging Centre (CUBRIC), Maindy Road, Cardiff, UK. CF24 4HQ. Email: steventonjj@cardiff.ac.uk Telephone: +44 (0) 29 206 88758*

(d) **Word count:** 2,994

(e) **Running title:** *An altered exercise response in HD*

(f) **Keywords:** *Huntington's disease, exercise, physiology, metabolism, cardiorespiratory*

2. Abstract

Background

Limited data suggests that an altered metabolic and cardiorespiratory exercise response may affect exercise performance in individuals with Huntington's disease (HD). There is no clear exploration of the response in individuals at different stages of the disease or in relation to genetic markers. This study aimed to examine the exercise response and recovery of HD participants, and the relationship to genetic and clinical markers.

Method

HD gene-positive participants (n=31; 9 pre-manifest; 22 manifest HD) and a healthy control group (n=29) performed an incremental exercise test until exhaustion. Performance, cardiorespiratory, metabolic and perceptual responses to exercise were determined from a maximal cycle ergometer test throughout the exercise test and during a recovery period.

Results

During sub-maximal exercise, metabolic (lactate levels, oxygen uptake) and cardiorespiratory markers (heart rate) were elevated in HD participants compared to controls. Lactate elevation was specific to pre-manifest HD participants. Work capacity was reduced in both pre-manifest and manifest HD participants with tests terminated with no difference in metabolic, perceptual or cardiorespiratory markers. Submaximal oxygen uptake was correlated with motor score, whilst peak measures were unrelated to genetic or clinical markers. Heart rate recovery was attenuated in pre-manifest and manifest HD participants.

Conclusions

Our findings confirm metabolic and cardiorespiratory deficits reduce exercise performance and affect recovery from an early stage in HD, with submaximal deficits related to phenotypic expression. Exercise capacity appears to be limited by an altered movement economy, thus clinicians should consider an altered exercise response and recovery may affect prescription in HD.

31 3. Introduction

32 Huntington's disease (HD) is a neurodegenerative disorder caused by the expansion of a polyglutamine
33 stretch within the Huntingtin gene¹. Neuropathology causes progressive motor disturbances,
34 cognitive dysfunction and psychiatric symptoms². However, as the mutant Huntingtin gene is
35 ubiquitously expressed, people with HD also experience an array of peripheral organ dysfunctions,
36 including a severe metabolic phenotype, weight loss, abnormal fat and glucose homeostasis,
37 cardiomyopathy, skeletal muscle wasting and reduced muscle strength³⁻⁶.

38 In healthy people, exercise interventions reduce all-cause mortality and are beneficial for
39 cardiorespiratory fitness, muscle strength, mental health and cognition⁷⁻¹⁰. Exercise prescription has
40 also been shown to be safe, feasible and in some cases, beneficial for motor symptoms in HD¹¹⁻¹³.
41 However, results have been variable and detrimental effects of exercise training have also been
42 reported^{4,14}. These conflicting and variable findings may reflect an altered physiological and
43 perceptual response to exercise in HD¹⁵ which may consequentially affect exercise prescription
44 outcomes. Importantly, the variability in the response to exercise suggests that targeted exercise
45 prescription may be required for people with HD.

46 In order to develop targeted approaches, it is necessary to understand what factors affect the
47 pathophysiological response to exercise. The current study was designed to improve knowledge on
48 who may benefit from exercise prescription by understanding individual exercise responses. Previous
49 work has found manifest HD participants experience an early increase in blood lactate and a reduced
50 work capacity and peak oxygen consumption, with no abnormalities in peak cardiac and ventilatory
51 performance¹⁶. Cardiac alterations and differences in perceived exertion were also found at lower
52 exercise intensities in HD participants prior to commencing an exercise intervention¹⁵, and an
53 increased oxygen cost of exercise has been demonstrated¹¹.

54 Previous studies in HD showing an altered exercise response were measured at a distinct
55 exercise stage, most commonly the response to maximal exercise intensity^{11,17}, whilst
56 cardiorespiratory and metabolic recovery after exercise has not been previously studied in HD. Here,
57 we measured the incremental response to exercise across the entirety of a cardiopulmonary test and
58 recovery period. We hypothesised that the physiological responses to the submaximal, maximal and
59 recovery phases may differ between controls and HD participants, which may relate to clinical and
60 genetic disease markers.

61 4. Methods

62 **Participants**

63 Thirty-one HD participants (22 manifest, 9 pre-manifest) were recruited from specialist HD clinics in
64 Cardiff, UK and Oxford, UK. Twenty-nine gene-negative controls were matched for age, gender and
65 self-reported physical activity levels. Demographic and clinical data are shown in Table 1. Data was
66 collected with ethical approval and participants gave informed consent.

67 **Exercise protocol**

68 Participants completed an incremental cycle ergometer exercise test¹⁸ (Excalibur Sport, Lode,
69 Netherlands) involving two minutes of rest, two minutes of unloaded cycling, followed by 25-watt
70 increments every two minutes, starting at 50-watts (Supplementary Figure 1). The exercise test was
71 symptom limited; individuals pedalled at 50 revolutions/minute until discomfort or fatigue due to
72 effort prevented them from maintaining the work rate. A 10-point Borg Scale measured perceived
73 exertion, fatigue in the legs and dyspnea. At 2 minute intervals, Borg ratings were collected along with
74 whole blood capillary lactate concentration (Lactate pro, UK).

75 Pulmonary gas exchange was measured on a breath-by-breath basis (MetaMax 3B, Cortex
76 Biophysik GmbH, Leipzig, Germany) and averaged every 30-seconds. Direct measurements of oxygen
77 consumption (O_2), carbon dioxide production (CO_2), minute ventilation (VE), and derived variables
78 including the respiratory exchange ratio (RER, i.e., O_2/CO_2), oxygen pulse (O_2/HR) and the ventilatory
79 equivalents for oxygen (VE/O_2) and carbon dioxide (VE/CO_2) were obtained. Heart rate was
80 continuously recorded using short-range telemetry (Polar S810, Finland).

81 After reaching exhaustion, participants cycled at an unrestrained speed at 25-watts for 2
82 minutes. Participants then transferred to a seated position and heart rate, blood pressure, lactate and
83 Borg ratings were recorded every 2-minutes for 10-minutes. Additional protocol details can be found
84 in the supplementary information.

85 **Clinical and genetic measures**

86 The motor component (TMS) of the UHDRS and the Total Functional Capacity (TFC)¹⁹ score were
87 recorded. Self-reported physical activity levels were recorded using the International Physical Activity
88 Questionnaire [IPAQ]²⁰. CAG repeat length data was available for 19 HD participants. To assess

89 baseline cognition, participants completed the Trail Making Test (part B) and The Symbol Digit
90 Modalities Test (SDMT).

91 **Statistical analyses**

92 HD participants were stratified according to disease stage (see Table 1; manifest HD participants = TFC
93 score < 13 and TMS >18, Supplementary Figure 2 shows TFC range in manifest group).

94 As sample size differed across test stages due to different termination points, multiple univariate
95 analyses of variance (ANOVA's) assessed group differences at each test stage (rest, submaximal stages
96 [50-, 75-, 100- and 125-watts], and peak performance), using SPSS software [IBM, version 23]. Multiple
97 comparisons were corrected for using the false discovery rate (FDR) at $p < 0.05$. Where results survived
98 the FDR correction, Bonferroni-adjusted post-hoc pairwise analyses examined group differences.
99 Multivariate ANOVA's were conducted for the recovery measures with post-hoc Bonferroni-adjusted
100 analyses.

101 Where group differences were found, the relationship with clinical and genetic data was tested using
102 a Pearson's correlation analysis.

103 A slope analysis assessed the relationship between physiology measures and work rate, from rest to
104 peak. The oxygen uptake efficiency slope (OUES)²¹, was determined from rest to peak using the
105 following equation:

$$106 \quad \dot{V}O_2 = a \log \dot{V}E + b;$$

107 *Where a represents the OUES, the rate of increase in $\dot{V}O_2$ in response to $\dot{V}E$.*

108

109 *[Table 1 here]*

110 **5. Results**

111 Despite no significant difference in age, BMI, and IPAQ score between control and HD participants
112 overall (all $p > 0.05$), manifest HD participants were significantly older than pre-manifest HD and
113 controls participants, thus age was included as a covariate in all analyses.

114 **Exercise protocol compliance**

115 Four HD participants were excluded from the analysis as they were unable to achieve the required
116 speed of 50 revolutions/min (see Supplementary Table 1). One HD participant experienced syncope
117 following maximal exertion.

118 **Resting measures**

119 Resting VO₂, heart rate, RER, mean arterial blood pressure (MAP) and lactate levels did not differ
120 between HD participants and controls ($p > 0.05$, Figure 1). Perceptually, there was no difference in
121 resting ratings of fatigue ($p > 0.05$).

122 **Submaximal exercise function**

123 Exercise parameters across test stages are shown in Figure 1.

124 **Physiology.** VO₂ was elevated in manifest HD participants compared to controls at 50 watts (33%
125 increase, $F_{2,50} = 5.42$, $p = 0.006$) and 75 watts (24.3% increase, $F_{2,50} = 3.45$, $p = 0.035$) with no
126 differences between pre-manifest and control participants ($p > 0.05$). Similarly, heart rate was
127 elevated in manifest HD participants compared to controls at 50 and 75 watts ($F_{2,49} = 5.54$ and 6.47
128 respectively, $p < 0.05$). In pre-manifest participants, heart rate was elevated compared to controls at
129 75 watts only ($p = 0.024$).

130 There was a main effect of group (HD vs. controls) on lactate levels at 50 watts ($F_{1,37} = 5.36$, $p = 0.035$),
131 75 watts ($F_{1,35} = 8.21$, $p = 0.028$) and 100 watts ($F_{1,34} = 5.90$, $p = 0.034$). Lactate was higher in pre-
132 manifest HD participants compared to controls, surviving Bonferroni correction at 75 and 100 watts (p
133 $= 0.014$ and 0.039) but not 50 watts (controls vs. manifest HD, $p = 0.053$). The difference between
134 controls and manifest HD participants was significant at 50 and 75 watts ($p = 0.027$ and 0.049
135 respectively) before age was accounted for, however with age as a covariate, this difference was not
136 significant.

137 Results from the slope analysis are shown in Supplementary Table 2. There was no difference in the
138 OUES nor the relationship between work rate and VO₂, heart rate and lactate between control and HD
139 participants (all $p > 0.05$).

140 **Perceptual responses.** There were no group difference in perceived exertion ratings at any workload
141 between 50 and 125 watts ($p > 0.05$).

142 *[Figure 1 here]*

143

144

145 **Peak exercise capacity**

146 Figure 2 shows the individual and group data at exhaustion. Age-adjusted marginal means are shown
147 in Supplementary Table 3.

148 **Performance.** Volitional exhaustion occurred at a lower working capacity (W_{peak}) in HD participants
149 ($F_{1,53} = 11.26$, $p = 0.001$, Figure 2A). Pre-manifest and manifest participants had a lower W_{peak}
150 compared to controls ($p = 0.049$ and 0.026 respectively); there was no difference in W_{peak} between
151 pre-manifest and manifest HD participants ($p > 0.05$).

152 **Physiology.** There were no differences in $\dot{V}O_2$ peak, heart rate, oxygen pulse, RER, VE, VE/VCO₂
153 (controls = 28.9 ± 0.8 ; pre-manifest = 30.7 ± 1.4 ; manifest HD = 30.0 ± 1.0), VE/O₂ (controls = $35.5 \pm$
154 1.4 ; pre-manifest = 33.2 ± 2.1 ; manifest HD = 34.9 ± 1.6) at exhaustion between HD and control
155 participants ($p > 0.05$).

156 Blood lactate production was 25.3% lower in HD participants at exhaustion compared to controls ($F_{1,40}$
157 = 9.37 , $p = 0.004$, Figure 2E) and was lower in manifest HD participants compared to controls ($p =$
158 0.032). Pre-manifest HD participants did not differ from controls. To determine the effect of W_{peak}
159 on lactate production, peak lactate levels were normalised by W_{peak} . There was no difference in
160 normalised peak lactate between control and HD participants (controls = 0.057 ± 0.003 ; pre-manifest
161 HD = 0.060 ± 0.004 ; manifest HD = 0.050 ± 0.003 , $p > 0.05$).

162 **Perceptual responses.** There were no differences in perceived exertion ratings between HD
163 participants and control participants at volitional exhaustion ($p > 0.05$).

164 *[Figure 2 here]*

165 **Exercise recovery**

166 **Physiology.** On a group level, heart rate did not differ between HD and control participants (Figure
167 3A). To examine individual variability in heart rate recovery (HRR), the change in heart rate from peak
168 was calculated. HRR was slower in HD participants compared to controls ($F_{10,70} = 1.98$, $p < 0.05$, Figure
169 3B). HRR was higher in controls compared to manifest HD participants at 2-minutes post exercise ($p =$
170 0.011), whereas after 4-minutes, HRR was higher in controls compared to pre-manifest and manifest
171 participants ($p = 0.038$ and 0.007 respectively). At 6-, and 8-minutes post exercise, HRR remained
172 higher in controls compared to pre-manifest participants ($p = 0.016$ and 0.021 respectively). There
173 were no differences in HRR between manifest and pre-manifest HD participants. Ten minutes

174 following exercise, the difference between HRR in pre-manifest participants and controls was not
175 significant ($p = 0.07$).

176 Lactate levels were reduced in manifest HD participants compared to controls at 2- and 4-minutes
177 post exercise ($p = 0.012$ and 0.034 respectively, Figure 3C), with no differences between control and
178 pre-manifest HD participants, and pre-manifest and manifest HD participants. To account for peak
179 lactate levels, lactate change (from peak) was calculated. There were no differences between control
180 and HD participants in lactate change during recovery ($p > 0.05$, Figure 3D).

181 **Perceptual response.** Self-reported ratings of fatigue during the recovery phase did not differ
182 between HD and control participants (Figure 3, $p > 0.05$).

183 *[Figure 3 here]*

184

185 **Relationship between exercise measures and genetic and phenotypic HD expression**

186 UHDRS TMS, where a higher score indicates greater movement disorder, was correlated with VO_2 at
187 50 watts ($r = 0.49$, $p = 0.032$), and 75 watts ($r = 0.44$, $p = 0.041$, Supplementary Figure 3). There was no
188 correlation between submaximal VO_2 and UHDRS TFC or CAG repeat length (all $p > 0.05$). Heart rate
189 at 50- and 75-watts, and lactate levels at 50-, 75- and 100-watts were not correlated with cUHDRS
190 TMS and TFC or CAG repeat length, all $p > 0.05$.

191 No correlations were found between clinical data (UHDRS TMS and TFC), CAG repeat length and peak
192 lactate or W_{peak} , all $p > 0.05$.

193 The correlation between HRR at 4- and 6-minutes post exercise and the UHDRS TMS ($r = -0.40$ and -
194 0.40 , $p = 0.039$ and 0.041 uncorrected) and UHDRS TFC ($r = 0.38$ and 0.46 , $p = 0.053$ and 0.017
195 uncorrected) was not significant after multiple comparison adjustments.

196

197 **6. Discussion**

198 This study shows an altered exercise response in people with HD during a graded exercise test. Despite
199 normal resting physiology, upon initiation of low intensity exercise, heart rate, lactate, and oxygen
200 uptake were elevated in HD participants compared to controls, suggestive of a movement economy
201 deficit and altered metabolism. The early elevation in lactate was observed in pre-manifest HD

202 participants specifically, suggesting that abnormal oxidative metabolism in skeletal muscle may be an
203 early feature of HD prior to functional decline. In contrast, elevated oxygen uptake at low intensities
204 was related to greater motor dysfunction in HD participants. In agreement with previous work¹⁶,
205 volitional exhaustion occurred at a lower working capacity in HD participants and both pre-manifest
206 and manifest HD participants produced less lactate at maximal effort. Almost one fifth of manifest HD
207 participants were unable to maintain the required cycling speed, highlighting an altered exercise
208 tolerance and physical ability in HD.

209 The altered cardiorespiratory and metabolic response to exercise may account for the variable
210 responses to exercise prescription previously reported in HD¹⁷. The relationship between oxygen
211 uptake during submaximal exercise and motor functioning suggest some of the variability can be
212 accounted for by phenotypic disease expression, however metabolic differences were not accounted
213 for by genetic load or phenotype, and were evident in pre-manifest HD participants, suggesting a
214 complex physiological response to exercise in HD.

215 Notably, these novel findings provide support for individualised exercise prescription, and
216 suggest that the effect of long-term exercise training may vary between individuals based on the
217 prescribed exercise intensity and their subsequent physiological response. Further studies measuring
218 the response to exercise are therefore recommended throughout the prescription period in order to
219 understand if disturbances in exercise response interact with exercise prescription outcomes. In
220 addition, it would be valuable to measure creatine kinase following exercise to explore muscle cell
221 disturbance in HD and to further explore autonomic functioning.

222 Whereas previous work has shown a reduced $\dot{V}O_2$ peak in manifest HD¹⁶, we failed to replicate
223 this. The discrepancy is most likely explained by the inclusion of age as a covariate in this study; when
224 VO_2 was measured as a percent of theoretical maximum capacity according to age, body type, and sex,
225 differences due to HD were no longer observed¹⁶. In line with previous work¹⁶, HD participants had a
226 normal ventilatory response to maximal exercise and both clinical data and genetic data did not
227 predict peak measures. CAG repeat length was also not predictive of the submaximal physiological
228 response, suggesting that genetic markers are insensitive to exercise response in HD. The rate of $\dot{V}O_2$
229 may be affected by how effectively oxygen is delivered to the muscles (respiratory and cardiovascular
230 systems), how it is utilised (metabolic systems) and how hard the muscles are working (neuromuscular
231 systems). We propose that higher VO_2 at submaximal exercise and not at peak may be due to muscles
232 working harder in people with HD due to chorea and/or poor co-ordination which requires greater
233 oxygen use. At peak exercise, aerobic metabolism is limited due to issues with bioenergetics and
234 mitochondrial functioning in the HD group.

235 After reaching exhaustion, heart rate recovery was attenuated in HD participants. This is a
236 novel finding; impaired heart rate recovery is an independent predictor of mortality in healthy
237 populations²². The fall in heart rate immediately after exercise is a function of the reactivation of the
238 parasympathetic nervous system, mediated by vagal reactivation in the first two minutes of
239 recovery²³. It is not clear whether vagal reactivation continues to mediate heart rate recovery for the
240 eight minutes observed here, or whether other mechanisms drive the slowed response. Due to the
241 cool-down protocol used, it is not possible to compare heart rate recovery with normative data
242 attained immediately after exhaustive exercise cessation, however our findings complement work
243 showing reduced cardiovagal modulation in middle-stage HD^{24,25}. The elevated heart rate observed
244 during submaximal exercise in HD participants replicates Dawes et al.¹⁵ and increases the robustness
245 of the observation that people with HD do not reach steady heart rates when initiating low-intensity
246 exercise. This is informative for exercise prescription; applying objective submaximal markers as
247 opposed to training intensities based on maximal measures (e.g. heart rate, VO₂), and measuring
248 exercise recovery as an indicator of intervention effectiveness may be more appropriate.

249 The difference observed in peak lactate is most likely driven by differences in task
250 performance. Working capacity was reduced in HD participants, and when peak lactate was
251 normalised for maximal work rate, the lactate differences were no longer significant. This suggests
252 that HD participants were unable to work as hard on the exercise test which contributed to the lower
253 lactate levels and is supported by previous work showing a reduced work capacity in manifest
254 participants¹⁶. The reduced working capacity in pre-manifest HD participants suggests an underlying
255 energy deficit rather than reduced muscle bulk.

256 Altered metabolism was also evident during submaximal exercise. The failure of oxidative
257 mechanisms can affect lactate production and clearance and previous work has shown elevated
258 lactate production in symptomatic HD participants during a cardiopulmonary test at 50 watts and 75
259 watts¹⁶. We replicated this finding, however, lactate was also significantly higher in pre-manifest HD
260 participants compare to controls at 75 and 100 watts, whereas the difference between controls and
261 manifest HD participants was not significant after accounting for age. The discrepancy in the results
262 may be partly explained by higher fitness levels in the current cohort and a broader symptomatic HD
263 group.

264 A common limitation in clinical exercise research is the physiological effect of medication.
265 Participants in this study were prescribed a diverse range of medications which alter metabolic
266 pathways²⁶. It is not known if observed metabolic and cardiac deficits are a cause or consequence of
267 HD toxicity, and/or mediated by medication. A comprehensive large-scale study is required to unpick

268 medication effects, with the metabolic consequences of drugs in the same sub-class varying
269 substantially due to differences in receptor pharmacology. Metabolic deficits observed in animal
270 models and cell cultures of HD suggest that observed effects are not purely driven by medication^{27,28},
271 however they may account for some variability in exercise prescription responses.

272 Overall, we have demonstrated that metabolic and cardiorespiratory deficits contribute towards a
273 reduced exercise performance and affects recovery in HD. Autonomic dysfunction has been reported
274 in a variety of neurodegenerative dementias^{29,30} and further work is required to elucidate whether
275 deficits are a direct consequence of peripherally expressed mutant huntingtin protein, or secondary
276 to a general decline in health or neurological dysfunction. This knowledge will be crucial for targeted
277 exercise prescription in HD and for determining outcome measures.

278

279 Acknowledgments

280 We gratefully acknowledge the participants in this trial, as well as their family members and
281 caregivers. We would also like to acknowledge the Waterloo Foundation, the Gossweiler Foundation,
282 the Wellcome Trust (200804/Z/16/Z), the Neuroscience and Mental Health Research Institute (Cardiff
283 University, UK); the NIHR Oxford Biomedical Research Centre (BRC) (UK) and the Oxford Health BRC
284 for funding this work.

285

286 7. References

287

- 288 1. Gusella, J. F., MacDonald, M. E., Ambrose, C. M. & Duyao, M. P. Molecular genetics of
289 Huntington's disease. *Arch. Neurol.* **50**, 1157–63 (1993).
- 290 2. Bates, G., Harper, P. S. & Jones, L. *Huntington's disease*. (New York: Oxford University Press,
291 2002).
- 292 3. Busse, M. E., Hughes, G., Wiles, C. M. & Rosser, A. E. Use of hand-held dynamometry in the
293 evaluation of lower limb muscle strength in people with Huntington's disease. *J. Neurol.* **255**,
294 1534–1540 (2008).
- 295 4. Kosinski, C. M. *et al.* Myopathy as a first symptom of Huntington's disease in a Marathon
296 runner. *Mov. Disord.* **22**, 1637–1640 (2007).
- 297 5. Lodi, R. *et al.* Abnormal in vivo skeletal muscle energy metabolism in Huntington's disease
298 and dentatorubropallidolusian atrophy. *Ann. Neurol.* **48**, 72–6 (2000).
- 299 6. Djoussé, L. *et al.* Weight loss in early stage of Huntington's disease. *Neurology* **59**, 1325–30
300 (2002).
- 301 7. Colcombe, S. & Kramer, A. F. FITNESS EFFECTS ON THE COGNITIVE FUNCTION OF OLDER
302 ADULTS: A Meta-Analytic Study. **14**, (2003).
- 303 8. Cooney, G. M. *et al.* Exercise for depression. *Cochrane database Syst. Rev.* CD004366 (2013).

- 304 doi:10.1002/14651858.CD004366.pub6
- 305 9. Arem, H. *et al.* Leisure Time Physical Activity and Mortality. *JAMA Intern. Med.* **175**, 959
306 (2015).
- 307 10. King, N. A., Hopkins, M., Caudwell, P., Stubbs, R. J. & Blundell, J. E. Beneficial effects of
308 exercise: shifting the focus from body weight to other markers of health. *Br. J. Sports Med.*
309 **43**, 924–927 (2009).
- 310 11. Frese, S. *et al.* Exercise effects in Huntington disease. *J. Neurol.* (2016). doi:10.1007/s00415-
311 016-8310-1
- 312 12. Khalil, H. *et al.* What effect does a structured home-based exercise programme have on
313 people with Huntington’s disease? A randomized, controlled pilot study. *Clin Rehabil* **27**, 646–
314 658 (2013).
- 315 13. Quinn, L. *et al.* A randomized, controlled trial of a multi-modal exercise intervention in
316 Huntington’s disease. *Parkinsonism Relat. Disord.* **31**, 46–52 (2016).
- 317 14. Potter, M. C., Yuan, C., Ottenritter, C., Mughal, M. & van Praag, H. Exercise is not beneficial
318 and may accelerate symptom onset in a mouse model of Huntington’s disease. *PLoS Curr.* **2**,
319 RRN1201 (2010).
- 320 15. Dawes, H. *et al.* Exercise testing and training in people with Huntington’s disease. *Clin.*
321 *Rehabil.* **29**, 196–206 (2015).
- 322 16. Ciammola, A. *et al.* Low anaerobic threshold and increased skeletal muscle lactate production
323 in subjects with Huntington’s disease. *Mov. Disord.* **26**, 130–7 (2011).
- 324 17. Dawes, H. *et al.* Exercise testing and training in people with Huntington’s disease. *Clin.*
325 *Rehabil.* 0269215514540921- (2014). doi:10.1177/0269215514540921
- 326 18. Collett, J. *et al.* Exercise for multiple sclerosis: a single-blind randomized trial comparing three
327 exercise intensities. *Mult. Scler. J.* **17**, 594–603 (2011).
- 328 19. Shoulson, I. & Fahn, S. Huntington disease: clinical care and evaluation. *Neurology* **29**, 1–3
329 (1979).
- 330 20. Craig, C. L. *et al.* International physical activity questionnaire: 12-country reliability and
331 validity. *Med. Sci. Sports Exerc.* **35**, 1381–95 (2003).
- 332 21. Baba, R. *et al.* Oxygen uptake efficiency slope: a new index of cardiorespiratory functional
333 reserve derived from the relation between oxygen uptake and minute ventilation during
334 incremental exercise. *J. Am. Coll. Cardiol.* **28**, 1567–72 (1996).
- 335 22. Cole, C. R., Blackstone, E. H., Pashkow, F. J., Snader, C. E. & Lauer, M. S. Heart-Rate Recovery
336 Immediately after Exercise as a Predictor of Mortality. *N. Engl. J. Med.* **341**, 1351–1357
337 (1999).
- 338 23. Imai, K. *et al.* Vagally mediated heart rate recovery after exercise is accelerated in athletes
339 but blunted in patients with chronic heart failure. *J. Am. Coll. Cardiol.* **24**, 1529–35 (1994).
- 340 24. Andrich, J. *et al.* Autonomic nervous system function in Huntington’s disease. *J. Neurol.*
341 *Neurosurg. Psychiatry* **72**, 726–31 (2002).
- 342 25. Bar, K. J. *et al.* Cardiovagal modulation upon postural change is altered in Huntingtons
343 disease. *Eur. J. Neurol.* **15**, 869–871 (2008).
- 344 26. Reynolds, G. P. & Kirk, S. L. Metabolic side effects of antipsychotic drug treatment –

- 345 pharmacological mechanisms. *Pharmacol. Ther.* **125**, 169–179 (2010).
- 346 27. Tsang, T. M. *et al.* Metabolic Characterization of the R6/2 Transgenic Mouse Model of
347 Huntington’s Disease by High-Resolution MAS ¹ H NMR Spectroscopy. *J. Proteome Res.* **5**,
348 483–492 (2006).
- 349 28. Naia, L. *et al.* Insulin and IGF-1 regularize energy metabolites in neural cells expressing full-
350 length mutant huntingtin. *Neuropeptides* **58**, 73–81 (2016).
- 351 29. Goldstein, D. S. Dysautonomia in Parkinson’s disease: neurocardiological abnormalities.
352 *Lancet. Neurol.* **2**, 669–76 (2003).
- 353 30. Chaudhuri, K. R. Autonomic dysfunction in movement disorders. *Curr. Opin. Neurol.* **14**, 505–
354 11 (2001).

355

356

357

358 8. Figure Legends

359

360 *Figure 1 Response to submaximal exercise. Results are marginal means adjusted for age, error bars = standard errors of the*
361 *mean. RER: Respiratory exchange rate. RPE: Ratings of perceived exertion. * p < 0.05 Bonferroni-adjusted. † unadjusted p <*
362 *0.05. preHD: pre-manifest, HD: manifest HD.*

363

364

365 *Figure 2 Peak exercise response at test termination. Black line represents mean; grey box depicts 95% CI. RER: respiratory*
366 *exchange ratio (VCO₂: VO₂); RPE: Rating of perceived exertion. PreHD: pre-manifest HD participants. HD: Manifest HD*
367 *participants. P-values are Bonferroni- adjusted.*

368

369 *Figure 3 Recovery response [B] & [D] show absolute change from peak. Results are marginal means adjusted for age. Error*
370 *bars = standard errors of the mean. RPE: rating of perceived exertion. HR: heart rate (beats/minute). * p < 0.05, ** p < 0.01.*