



Neonatal nurses' response to a hypothetical premature birth situation: What if it was my baby?

Janet Green

University of Technology Sydney, Australia

Philip Darbyshire

Monash University, Australia; Flinders University, Australia

Anne Adams

Retired, Australia

Debra Jackson

Oxford Brookes University, UK; Oxford University Hospitals NHS Foundation Trust, UK

Abstract

Background: Evolving technology and scientific advancement have increased the chances of survival of the extremely premature baby; however, such survival can be associated with some severe long-term morbidities.

Research question: The research investigates the caregiving and ethical dilemmas faced by neonatal nurses when caring for extremely premature babies (defined as ≤ 24 weeks' gestation). This article explores the issues arising for neonatal nurses when they considered the philosophical question of 'what if it was me and my baby', or what they believed they would do in the hypothetical situation of going into premature labour and delivering an extremely premature baby.

Participants: Data were collected via a questionnaire to Australian neonatal nurses and semi-structured interviews with 24 neonatal nurses in New South Wales, Australia.

Ethical considerations: Relevant ethical approvals have been obtained by the researchers.

Findings: A qualitative approach was used to analyse the data. The theme 'imagined futures' was generated which comprised three sub-themes: 'choice is important', 'not subjecting their own baby to treatment' and 'nurses and outcome predictions'. The results offer an important and unique understanding into the perceptions of nursing staff who care for extremely premature babies and their family, see them go home and witness their evolving outcomes. The results show that previous clinical and personal experiences led the nurses in the study to choose to have the belief that if in a similar situation, they would choose not to have their own baby resuscitated and subjected to the very treatment that they provide to other babies.

Conclusion: The theme 'imagined futures' offers an overall understanding of how neonatal nurses imagine what the life of the extremely premature baby and his or her family will be like after discharge from neonatal

intensive care. The nurses' past experience has led them to believe that they would not want this life for themselves and their baby, if they were to deliver at 24 weeks' gestation or less.

Keywords

Ethics, extreme prematurity, hypothetical, neonatal nurses, qualitative research

Introduction

There is no argument that the survival of extremely premature babies has improved substantially; however, similar improvements have not been noted for long-term morbidity¹ or in the long-term problems associated with being born at an extremely early gestation. The minimum age of viability is considered to be as young as 23 weeks' gestation, with the occasional survivor reported at 22 weeks' gestation.² Neonatal clinicians cannot definitively predict the individual prognosis of an extremely premature baby,³ but it becomes problematic when they cannot predict if survivors will have a chronic or disabling condition.⁴ Unsurprisingly, the lower the gestational age, the higher the mortality and morbidity¹ and the greater the chance of long-term and severe morbidities.⁵

The current outcomes for extremely premature babies related to major neurological impairments have remained stable, despite improved survival.⁶ Such neurological damage may include brain injury with subsequent physical and cognitive disability resulting from intraventricular haemorrhage and periventricular leukomalacia.⁷ Other potential problems include chronic lung disease, hearing impairment, sight problems caused by retinopathy of prematurity and short gut syndrome from necrotising enterocolitis.^{8,9} For approximately half of the survivors of extreme prematurity, special education services will be needed.¹⁰ The outcomes of extreme prematurity reported in even recent literature published yesterday does not necessarily reflect the outcome that would be reported today because cohorts may have been from the last 10 years, and it might have taken 2 years for the research to be published. However, it is the best that is available at this point in time. Treatment regimes constantly change, making it difficult to interpret outcomes research. While it is clear that extremely premature babies can develop into healthy, independent children and adults,¹¹ this is not always the case, and truly 'informed' consent would dictate that any discussions with parents should consider the possibility that an extremely premature baby may survive with a major disability.

A hypothetical situation is one involving or based on a suggested idea or theory that invites us to consider 'What if...?' or 'Imagine that...?'. It involves or is based on a hypothesis with a not real or imagined example. Words associated with hypothetical include suppositional, theoretical and speculative.¹² Hypothetical situations are useful because it is often necessary to hypothesise in order to explain how people believe they will behave or act in certain situations. Knowing what one would do in a hypothetical situation is often more than an educated guess, and in fact, hypothetical scenarios may not accurately reflect actual behaviour because individuals can have difficulty projecting themselves into the future with accuracy.¹³ Human beings weigh up evidence and many other factors when coming to a decision about a particular issue. For neonatal nurses, contemplating the outcome for babies of 24 weeks' gestation and less, and the perceived burden on their family, could make them consider the hypothetical situation of 'What if it was me and my baby'?

Research design

Aim

The findings presented in this article are part of a larger mixed-method doctoral thesis¹⁴ that explored the ethical issues and caregiving experiences of Australian neonatal nurses who cared for extremely premature babies of 24 weeks' gestation and less. Other publications from this research include the difficulties

associated with caring for extremely premature babies who at times look more like a foetus than a human baby;¹⁵ the burden of keeping secrets and how neonatal nurses have to keep information to themselves.¹⁶ Three further papers include caring for parents whose belief in a miracle was all consuming,¹⁷ how neonatal nurses manage the uncertainty associated with caring for extremely premature babies¹⁸ and the difficulties neonatal nurses face in trying to reconcile hurting with caring, when they inflict pain on extremely premature babies as part of their treatment.¹⁹ The myth of the miracle baby has been challenged in another paper,²⁰ the anguish and desperation to become parents has been explored,²¹ as has the quality of life of extremely premature infant survivors.²²

The focus of this article is the quantitative and qualitative data that explore the neonatal nurses' perception of why they feel that, hypothetically, they would not want their own extremely premature baby resuscitated and treated if they found themselves in the same position as the parents of an extremely premature infant.

Data collection

Two data collection methods were used in this research. In the first stage of the study, Australian neonatal nurses ($n = 414$) were surveyed using a self-completed questionnaire that was analysed using SPSS. In the second stage, purposive sampling was used and data collected through 14 semi-structured interviews with 24 neonatal nurses from the state of New South Wales (NSW) and the Australian Capital Territory (ACT).

The questionnaire was a 64-point Likert, paper-based survey that sought to explore the attitudes and the legal, ethical, social and technological issues experienced by neonatal nurses when providing care to extremely premature babies. The questionnaire was modified with permission from a study undertaken by Armentrout.²³ Questions were based on a literature review that determined current issues and concerns surrounding the provision of care to extremely premature babies. The questionnaire addressed neonatal nurses' demographics, age and years of experience in caring for extremely premature babies. Both open and closed questions were included, with opportunities for participants to comment on their experience and issues of concern about caring for extremely premature babies.

In all, 14 interviews were conducted involving 24 participants. There were eight single interviews and six focus groups comprising between two and six neonatal nurses. All data were collected by the first author, an experienced neonatal nurse. The interview questions explored the nurses' experiences of caring for babies of ≤ 24 weeks' gestation. The interview questions were constructed from the significant issues arising from the questionnaire, content analysis of the open-ended philosophical questions from the questionnaire and other issues that emerged during the interviews.

The interviews were semi-structured; however, time was also allowed for unstructured conversation. The interviews occurred in different locations; the interviewer's home (4), in the participants' own homes (5) or a quiet room at the participant's hospital of employment (5). The duration of the interview was between 60 and 90 min. The full interviews were transcribed prior to in-depth analysis to identify major themes. This article draws on the questionnaire data, the qualitative data from the questionnaire and the interviews.

Setting and participants

Participants who were interviewed included Registered Nurses who were currently employed in a neonatal intensive care unit (NICU) where extremely premature babies are cared for, a paediatric intensive care unit where babies are cared for, or members of the newborn emergency retrieval team. The nurses required greater than 5 years' experience in caring for babies ≤ 24 weeks' gestation. They needed to be English speaking, willing to participate in the research and to agree to have the interview recorded. The extremely premature baby, especially in the first 24 h, can be fragile and critically ill, thus requiring the most experienced and skilled nursing staff to care for them. These babies are not given to inexperienced nurses

to care for; therefore, those nurses with 5 years or more experience with caring for extremely premature babies would have the experience required to be interviewed. All 24 of the nurses interviewed were females, 10 were mothers, 14 were childless but within childbearing age (33–45 years).

Ethical considerations

Questionnaire and interview participants were provided with a participant information sheet. Verbal and written consent was obtained from the interview participants, and they were given the option to ask questions for clarification or withdraw from the study. The confidentiality of all participants was assured, first by the anonymous procedures associated with the questionnaire. Second, the names of the interview participants were not included on the transcripts and third, the data were secured in a locked drawer. This research project was approved by the Flinders University of South Australia Social and Behavioural Research Ethics Committee (Approval Number 1924). The topic of this research is considered a sensitive one; therefore, counselling was made available to the interview participants if required, although none of the nurses required this service.

Data analysis

A qualitative method informed by phenomenological insights and the work of Van Manen²⁴ were considered the most appropriate way to interpret the interviews because of the need to understand the nurses' experiences of caregiving dilemmas surrounding extremely premature babies. Phenomenology is the study of lived experience and asks the question 'what is this or that kind of experience like'.²⁴ Van Manen's²⁴ approach is that reflecting on the lived experience cannot occur while the person is still living it. The researcher is required to capture the retrospective reflection of the nurse who has provided care to extremely premature babies.

The procedural component of phenomenology requires that the text from the interviews be examined carefully and systematically. Formal phenomenological analysis requires line-by-line analysis, and the discovery and construction of themes. Creating themes is an active interpretative process, in keeping with the work of Van Manen.²⁴ Creating themes helps the researcher identify the significant issues in the data. The search for the 'big picture' according to Braun and Clarke (p. 12)²⁵ means that the theme is authentic and provides an accurate understanding of the nurses' experience. Accounts of the nurses' experiences were structured to form the whole, or a full description of the phenomenon.

A qualitative study needs to be rigorous. The specifics of rigour is about how the research paradigm's ontology and epistemology inform the interpretative methodology in the study, and whether or not it is able to answer the question under investigation. The validity of a qualitative study is about trustworthiness, and how the researcher describes and interprets what has been told to them by the informant.²⁶ If the lived experience is captured, there will be richness in the description, which will lend credibility to the research and study. Readers also establish the trustworthiness if they consider the study is transferable to another context, and are able to follow the decision trail of the researcher. The researcher should be able to provide confirmability of the study, or how the qualitative interpretations came to fruition. All data sources were collected by the first author, and all data and interpretations were regularly audited and validated by the co-authors.

Findings

This article outlines the neonatal nurses' consideration of the hypothetical situation of what would they do if they delivered an extremely premature baby. It is important to emphasise that the results reported in this article are applicable only to those babies of 24 weeks' gestation and less, and are not generalisable to any other neonatal population, as the outcomes for extreme prematurity have a greater complexity than the

outcomes for babies of other ages nursed in the NICU. Nor are the results readily generalisable to neonatal nurses other than those interviewed. As there are two data sources comprising the results, the qualitative results of the questionnaire are represented as (Q response number) and the interview transcripts are represented as (Nurse number).

Questionnaire data

There were no specific questions in the questionnaire that asked about what the nurses would do if they personally delivered an extremely premature baby; however, their experience has led them to believe they would not want their own baby to be resuscitated and treated. The questionnaire data are included to add context to the qualitative data. Only nurses with experience with caring for babies of 24 weeks' gestation and less were asked to complete the first part of the questionnaire (Tables 1 to 4).

Table 1. Results – caring for babies of 24 weeks' gestation and less makes me feel hopeful.

	Frequency	Percentage
Never	35	10.6
Seldom	120	36.5
Occasionally	147	44.6
Almost always	27	8.3
Total	329	100

Table 2. Results – caring for babies of 24 weeks' gestation and less makes me feel concerned.

	Frequency	Percentage
Never	1	.3
Seldom	6	1.8
Occasionally	35	10.6
Almost always	290	87.3
Total	332	100

Table 3. Results – caring for babies of 24 weeks' gestation and less makes me feel positive.

	Frequency	Percentage
Never	40	12.1
Seldom	136	41.1
Occasionally	130	39.3
Almost always	25	7.5
Total	331	100

Table 4. Results – caring for babies of 24 weeks' gestation and less makes me feel discouraged.

	Frequency	Percentage
Never	12	3.6
Seldom	43	13.1
Occasionally	194	58.9
Almost always	80	24.2
Total	329	100

The results show that nurses were rarely hopeful (44.6% occasionally hopeful and 36.5% seldom hopeful) about the survival and outcome for extremely premature babies.

The majority of nurses (87.3%) were concerned about the outcomes for the baby and parents.

The results show that nurses usually found it difficult to be positive (39.3% occasionally positive and 41.1% seldom positive) about the survival and outcome of extremely premature babies.

The results show that many nurses were discouraged (58.9% occasionally discouraged and 24.2% almost always discouraged) about the survival and outcome of extremely premature babies.

Interview data

The theme 'imagined futures' offers an overall understanding of how neonatal nurses imagine what the life of the baby and family will be like after discharge from the NICU. What the nurses had seen in the past had made them believe that they would not want this life for themselves and their baby if they were to deliver at 24 weeks' gestation or less. The theme of 'imagined futures' comprised three sub-themes: (1) 'choice is important', (2) 'not subjecting their own baby to treatment' and (3) 'nurses and outcome predictions'. The first theme 'choice is important' showed that the neonatal nurses believed that they should have the choice to decide whether their baby born at 24 weeks' gestation and less should be resuscitated. They also believed that the parents of other babies in the NICU should also have the same choice. The second theme 'not subjecting their own baby to treatment' was about the nurses imagining what might happen, and what they would do if they delivered an extremely premature baby. The third theme 'nurses and outcome predictions' explores how the nurses made their decisions about whether they believed a baby would have what they considered to be a good or poor outcome.

Choice is important. The nurses expressed that they would not have their own baby treated if they delivered at 24 weeks' gestation and less. This issue was noted in the questionnaire when approximately 25% of respondents wrote unsolicited comments, for example, 'every neonatal nurse I know would go bush rather than go to a tertiary centre if they were in prem labour with a pregnancy of 24 weeks gestation or less' (Q response 106). In colloquial Australian language, the words 'go bush' are used to mean that in remote areas (the bush) there would be no access to a NICU, and hence no chance to save an extremely premature baby:

I am glad I had term babies. I would hate to have a 23–24 weeker and 'in theory' would actively discourage any member of my family from insisting on full resuscitation and treatment if they were unfortunate enough to go into preterm labour, or require delivery of a 23–24 weeker. As I said though that is the theory – I hope I would have the strength in practice. (Q response 199)

Only one of the 24 nurses interviewed stated she would have her baby treated, but the caveat was that only if treatment would be withdrawn if the baby suffered a large intracranial bleed. The nurses all considered treatment of the baby to be a personal autonomous choice. One nurse claimed her decision was an educated one, based on her understanding that an extremely premature baby breathing at birth did not equal sustained life. The decisions the nurses made were seen as recognition of the reality that parents faced during treatment. One nurse summed up the situation when she stated 'if you were given a 10% chance of a successful outcome from surgery, would you take the chance?' (Q response 368). It was the understanding of the nurses that the mere survival of the extremely premature baby did not constitute a positive outcome.

All respondents claimed that just as they had the right to choose, the parents had the same rights, but the ability to make decisions comes with information. It was accepted that parents might make decisions based on how the information was presented to them. One nurse explained the difficulty:

I don't know what they [Doctors] tell them. They must tell them survival. I don't think they tell them intactness [intact survival]. If someone told me that, 'Yes at 23 weeks you've got a 50% chance of survival, but a high chance of being abnormal', I don't think I'd want my baby resuscitated. (Nurse 19)

This was a hypothetical situation, and it is acknowledged that the nurses would not know in reality what they would do until they were in that situation. It was interesting that they wanted other people's babies to have the opportunity to live, if that is what they chose, yet not want that for their own child. One nurse wrote on her questionnaire,

if I had a baby 24 weeks or less, I would hope the doctor would listen to me about my wishes of not having the baby treated. This has always been my wish, and I do not intend to change it. (Q response 296)

Not subjecting their own baby to treatment. Knowledge about the outcomes of extremely premature babies was considered powerful knowledge by the nurses. They were generally emphatic that they would not consent to treatment if they were to deliver an extremely premature baby. There are always difficulties in talking about hypothetical situations. The nurses spoke as if they knew what they would do. They made it clear that if they started to labour, they would avoid any centre capable of providing treatment for an extremely premature baby. They feared having the decision to treat being taken out of their hands. Most nurses said they would present to a tertiary care facility only if they feared for their life. Two nurses in a focus group stated,

Unless I was bleeding and feared for my own life, I would sit at home. I would labour at home, deliver at home and then wrap it and just cover it until it died. Then present [to hospital] with the dead baby. I would not present in with something you could ventilate. (Nurse 19)

You could still present, it doesn't mean they have to do anything. (Nurse 20)

That would be my fear that it came out and it cried, and they did something and then there was no pulling out [withdrawing treatment]. And if it cried they would take it off me and they would bring it here [a NICU]. That would be my fear. (Nurse 19)

The nurses were convinced they had a realistic perception of extreme prematurity because of their experience in working with these babies every day and seeing babies come back who had a less than favourable outcome, and whose parents had told them of their day-to-day existence. When speaking of her hypothetical reluctance to have a baby of 24 weeks or less resuscitated, one nurse stated in a matter of fact way,

I wouldn't leave it for dead or anything, but the baby wouldn't survive. I'm afraid I'm not one of those courageous people who can see it as God's plan or anything else. (Nurse 14)

Caring for other people's extremely premature babies, but not wanting their own treated represented a contradiction and a possible conflict for the nurses. One nurse explained she would not present to hospital because of her belief that presentation implied she was seeking treatment. She stated, '... if I do present at the hospital 24 weeks pregnant, I'm asking for help. Usually the patient, they do want something done' (Nurse 11). In this situation, informed consent was needed and initial and ongoing consent is required for all treatment. The nurses might be alluding to the emotional aspects of seeking help for an extremely premature baby. Many nurses held that they would stay away so they could not be persuaded to have their baby treated, or have the staff treat the baby without consent. Several stated they had read the book, 'The Long Dying of Baby Andrew' by Peggy Stinson and Robert Stinson²⁷ and although this book is old it remains a 'must read' for every member of the neonatal team. The book is about treatment without consent and experimentation in

the name of progress. In this study, fear was a powerful motivator for these nurses. Fear of the technological imperative would keep them away from any centre capable of resuscitating and treating tiny babies. It was not the fear of resuscitation that drove these nurses to essentially avoid their own workplaces, it was their fear that treatment would not be withdrawn when they believed that their baby may have been irreparably damaged. Technology, they felt, seemed to take on a power of its own, and once instituted it could be exceptionally hard to withdraw. The nurses understood that some medical staff would not, '... pull the plug' (Nurse 16), and the nurses were not prepared to take the risk.

When the nurses considered what was required to keep an extremely premature baby alive, they were not surprised when some parents chose not to have their baby resuscitated. When asked if she would have her own baby resuscitated, one nurse replied,

I come back to the amount of intervention at a very basic sense of the word, and I intervene with my large plastic tube [ETT] to keep this child alive. I invade its body, I create holes in it to make this child live. And if I can give out the relationship on that small baby to the size of the plastic and try to say 'how big would that bit of plastic be in relation to me'. I find it completely unacceptable if I had to accept a UAC [umbilical artery catheter], a UVC, [umbilical vein catheter] a peripheral cannula, the size of the endotracheal tube, even a chest drain. All of these things in my baby. I know if I had the decision I would say definitely not. (Nurse 6)

Knowing what to do in hypothetical situations is difficult. It is even more difficult to know what neonatal nurses would do if they were in premature labour. Gaining an accurate picture of what nurses believe they would do in a hypothetical situation is difficult. Unless faced with the situation, it is still very much a nurse believing she would know what to do in a given situation. That said, what is interesting is that the cognitive dissonance of the nurses meant that they would avoid at all costs the very places where they work, where they practice their profession and where they provide care to 'other parents'.

Nurses and outcome predictions. Not every neonatal nurse is directly aware of the outcomes for individual extremely premature babies. The nurses in this study had all read literature on the outcomes for tiny babies and had seen babies return for follow-up. It was the opinion of the nurses that '... the literature ... it's still a bit weighted to make things look really good ... There would probably be hardly any babies that came out unscathed' (Nurse 19). Not all neonatal nurses are aware of the outcomes of extreme prematurity; however, the nurses in the study knew about the possible long-term outcomes. One nurse explained,

That's something that we can demonstrate by the amount of reading that we've done and the facts that we have gathered ... The experience, what we've seen, what [the babies and families] we have nursed. You know the outcomes that we've seen. (Nurse 1)

Many nurses spoke of how they heard about the outcomes of babies. Word filtered back to the NICU. Most nurses, however, focused on the poor outcomes:

We do hear of all the follow-ups and there are a lot of positive follow-ups as well. I guess we tend to hear more of the disasters. (Nurse 13)

The nurses did not often see the good outcomes in tiny babies. It was difficult because '... we hear more of them [poor outcomes] than the ones who do well. That's the problem isn't it, we hear about the ones who are disasters' (Nurse 14). In using everyday language, this nurse was showing concern about causing damage to the baby during treatment. It is hoped that this form of language would not be used outside the confines of nursing. The nurses spoke of what they call 'disasters' (Nurses 13, 14 and 19)

when referring to outcomes. A disaster implies devastation and distress and three nurses perceived some babies in this way.

The nurses all liked to hear about positive outcomes. One nurse explained,

Occasionally we get to see how babies are doing years down the track. That's positive. It's not that often, though.
(Nurse 14)

Several nurses spoke about their colleagues who were optimistic about extremely premature babies, but who were not familiar with outcomes nor had they attended the follow-up clinic. This situation was not uncommon, according to the nurses, because of the way that NICUs are structured, with an acute area separated from another area for less acute babies. Nurses who chose to work in intensive care might not get to see babies come back to the unit once they had been discharged. One nurse stated,

We see a small facet. We don't see it [the baby] often down the track, or it's walking around blind. We don't see those issues, and if you don't see something you don't realise. (Nurse 5)

Another nurse explained that when she undertook her neonatal nursing course, the consultant neonatologist was adamant the nurses should go to the follow-up clinic to see the babies they had cared for. Here they, '... saw the so called bad outcomes' (Nurse 6). This was beneficial for the nurses, one nurse stating, 'I found that extremely confronting in there... for a long time I'd always thought that the children who didn't survive intact were the mistakes' (Nurse 6). One nurse emphasised she knew about the poor outcomes, but chose to focus on the positive outcomes. This nurse explained, '... it's probably a survival mechanism. You switch them out of your brain as much as possible' (Nurse 24). Another nurse was convinced that constantly focusing on the negative outcomes made it difficult for her to continue working in the NICU. She stated, '... you just want to see that baby go home and not really think of what's going to happen later on' (Nurse 11). The nurses felt it was easier for them to concentrate on the positive aspects. Dwelling on the negative aspects might have been too confronting for them professionally.

All nurses interviewed believed they knew which babies would do well, and which babies would have poor outcome. They based their predictions on technical information and the baby's reaction to handling and its response to caregivers. Several nurses acknowledged they had been wrong in their predictions for extremely premature babies. The nurses all spoke of times when they believed a baby would not survive, and yet it survived. They emphasised that survival was not related to outcome. This means they might not be correct in predictions of survival, but were often correct in predictions about outcomes. One nurse explained,

When it comes back for follow up, you think 'Oh yeah you're not as good as what they hoped'. But the fact that you merely survived, is quite miraculous. (Nurse 12)

One nurse, who had seen two of these babies have good outcomes, still had a spark of hope:

I might think the child needs to be turned off [had treatment withdrawn]. I thought a couple of times babies... I really thought they should be turned off and they should die. They [medical staff] have persevered and these people [babies] have come back OK. (Nurse 24)

Discussion

A large body of literature indicates that children born extremely preterm are likely to experience ongoing problems; however, the extent of those problems continues to be debated, and there is much uncertainty.

How neonatal nurses manage the uncertainty of caring for extremely premature babies has been elucidated by Green et al.¹⁸ What is known is that a baby born at 24 weeks' gestation could be 650 g, have a head circumference of 22 cm and be 31 cm long;²⁸ therefore, it is likely to be smaller in all aspects of growth and have a poorer developmental outcome. Gestational age exerts the greatest influence on the survival²⁹ and outcome of premature birth.³⁰ The outcome literature reports on cohorts of very low birth weight (VLBW) infants (less than 1500 g), and extremely low birth weight (ELBW) infants (less than 1000 g); however, it is difficult to gain a full picture of the babies who were the 'microprems' or 24 weeks' gestation and less. Lower birth weights and gestation consistently have poorer outcomes,³¹ yet it is difficult for the authors of this article to specifically identify the population under consideration in the outcomes literature and make judgements about their outcome to see if the neonatal nurses are correct in their concerns about poor outcome, and why they would not want their own baby resuscitated and treated if they delivered at 24 weeks' gestation or less. Significantly perhaps, the outcomes literature reports de-identified cohorts rather than the individual babies that neonatal nurses see in their practice. Perhaps this might explain the discrepancy between the outcomes literature and the views that neonatal nurses express in the current research. Nurses' perceptions of outcomes and their sense of 'What they would do if this was their pregnancy and baby' may be more powerfully shaped by seeing an individual baby with a poor outcome going home with its family. What they have seen in the past has affected how they would envisage their future if they were to go into premature labour.

Making decisions about the resuscitation and treatment of extremely premature babies in industrialised nations are complex, and Australia is no exception. Australian guidelines recommend that from 20 to 22 weeks' gestation, life-sustaining treatment should usually not be provided. At 23–24 weeks' gestation, life-sustaining treatment should be guided by parental wishes, and from 25-week life-sustaining treatment should usually be provided.^{32,33} Although the nurses believed that having choices was important for parents of extremely premature babies, they knew from previous experience that such choices were not absolute, and medical staff could override parental decisions. Complexities exist, and Larcher³⁴ states that decisions made about babies at the limits of viability cannot be made on the basis of clinical facts alone, and should take into account the values and belief of all concerned, including the staff. However, differences in moral beliefs exist and can be compounded by a power imbalance inherent in the professional–parent relationship.³⁴ This creates a dilemma because 55% of neonatologists in a study by Weiss et al.³⁵ stated they would not resuscitate within the 22nd week of gestation, which could mean that 45% would resuscitate during the 22nd week. Neonatologists in the same study were less likely to consider the parent's opinions all the time when choosing whether or not to resuscitate.³⁵

A further complexity in the decision-making process is the religious beliefs of the attending neonatologist. In all, 89% of neonatologists in a study by Donohue et al.³⁶ reported their religious beliefs influenced their medical practice. Lawrence and Curlin³⁷ found that doctors with committed religious beliefs gave less weight to patient's expressed wishes. As Paris et al.³⁸ have lamented, physicians can resuscitate no matter how premature, how unlikely to survive, how likely to incur severe disabilities or how strongly the parents object, showing that parental choice is not absolute.

In decision research, the use of imagined or hypothetical situations is accepted as a legitimate way of studying real behaviour.³⁹ When making decisions based on a hypothetical, a subject considers events that may or may not happen, considers feelings that they may or may not have, while anticipating possible outcomes and evaluating those outcomes within preferred options.³⁹ In research that uses a hypothetical methodology, vignettes are used as the equivalent of a case study where respondents reveal their perceptions and values.¹³ In the current research, the nurses were not given a vignette, but rather they used their own experiences of caring for extremely premature babies and their families to make the decision that they would not have their baby of 24 weeks' gestation and less resuscitated and treated. These nurses created their own vignette.

It is quite reasonable that the nurses would project themselves into the future in this way. Extrapolation to the psychological literature shows that although the future is unknowable, humans are able to simulate potential future scenarios or mental representations of the future in their minds.⁴⁰ The ability to construct mental representations of episodic future thinking is an important cognitive event. The ability to remember past events and imagine future events relies on the storage and retrieval of information in the episodic memory.⁴⁰ Knowledge about the future is structured around personal goals and cultural life scripts which are the person's expectations about the order and timing of events in his or her life.⁴⁰

A critical factor to be considered in hypothetical situations is the importance of the decision because important issues are afforded more cognitive processing which leads to better and more deeply considered decisions.³⁹ Anticipated emotions such as disappointment and regret can be operative in decisions in both hypothetical and actual situations.³⁹ There are such things as hot and cold states when it comes to hypothetical decisions. A hot state is important and is when a person makes a hypothetical decision that would have important consequences if made for real.³⁹ The nurses in this study were projecting future outcomes. Therefore, they were ultimately considering the consequences of having their own baby given intensive care, and if irreversible damage occurred, it would have significant real-world consequences.

The neonatal nurses stated they would not want their babies born extremely prematurely to be treated. This is in conflict with Sanders et al.⁴¹ who found the opposite for neonatologists, albeit over 20 years ago. They found 61% of neonatologists would want their child treated with the same level of aggression with which they treated patients, while 34% would want their child treated less aggressively.⁴¹ Streiner et al.⁴² found physicians were more optimistic than neonatal nurses about the outcomes of extremely premature infants. In a large study that included 3425 nurses by De Leeuw et al.,⁴³ it was found that the nurses were more prone than doctors to withhold treatment on a 24-week gestation infant. Not surprisingly, given the importance of choice identified by the nurses in this study, that the nurses in De Leeuw et al.'s⁴³ study would ask parental opinion about further treatment choices.

In all, 88% of nurses and 85% of neonatologists in a study by Oei et al.⁴⁴ would almost always resuscitate 24-week gestation infants; however, only 24% of neonatologists and 6% of nurses would resuscitate a 23-week gestation infant. These clinicians were more optimistic about survival than the long-term outcome. Janvier et al.⁴⁵ authored a paper titled 'Nobody likes premies', where they gave a hypothetical scenario to physicians and students. The results make them question whether there is a 'systematic devaluation of the newborn due to deeper rooted anthropological, cultural, social and evolutionary factors'.⁴⁵ There is nothing in the current research that would support Janvier et al.'s⁴⁵ finding. McHaffie and Fowlie⁴⁶ found neonatal nurses would not want their own child to go through the painful therapies when the benefits were questionable. The outcomes for extremely premature babies were a similar concern to the neonatal nurses in McHaffie and Fowlie's⁴⁶ research.

Janvier et al.⁴⁷ asked resident doctors and nurses involved in perinatal care if they would resuscitate a 24-week gestation infant 'depressed' at birth. They gave a vignette that gave the outcome of extreme prematurity, but does such an outcome change if the baby needs significant resuscitation? In all, 21% said they would resuscitate. The authors wondered if it was the 'emotional response to the foetus of 24 weeks that looks barely human'.⁴⁷ The difficulties for neonatal nurses when caring for the extremely premature baby that does not have the widely accepted look and characteristics of normal 'babyness' have been discussed by Green et al.¹⁵ Interestingly, for this article, the foetal look of the baby had nothing to do with why the nurses would not want their own baby resuscitated. Janvier et al.⁴⁷ did not find that nurses were less interventionist than the resident physicians; however, they did find that the nurses' attitudes reflected their personal experiences of seeing many babies with complications. These nurses worked in an outborn surgical centre. The nurses in this study worked in a variety of areas including outborn surgical centres. Janvier et al.⁴⁸ in citing the same study that the 'more the major complications are seen, the more nurses have a false impression of bad outcomes'. Janvier et al.⁴⁷ suggest that the unwillingness to intervene for extremely

premature infants is out of proportion to the prognosis. Yet, Janvier et al.⁴⁷ created the scenario of a depressed baby, which means it needs resuscitation. The poor outcomes of cardiopulmonary resuscitation (CPR) in the premature infant lead Lantos et al.⁴⁹ as early as 1988 to believe that CPR is a non-validated therapy in extremely premature babies and that these infants should not be subject to a standing order for CPR in the first few days of life. They argue that aggressive support should still be given, but the need for CPR should be taken as a sign of impending death. Barr and Courtman⁵⁰ agree and suggest that CPR be withheld in extremely premature babies who do not have a reversible cause for their cardiac arrest.

McHaffie and Fowlie⁵¹ found neonatal nurses would want fewer interventions for their own extremely premature babies than that currently offered to patients in their NICU of employment. Perhaps their experience has led them to believe that not all babies can survive, and should it happen to them they would prefer not to be held 'hostage to circumstance'.⁵² Harrison⁵² spoke of a midwife who had contingency plans in case of premature delivery, and '... she would drive to a remote area, as far from a tertiary unit as possible, and let nature take its course'. The following is a quote from McHaffie and Fowlie⁵¹ where a neonatal nurse explains her beliefs:

I always try to think, if it was my baby, what would I want, and then weigh that up against what I know as a professional ... if I delivered a 23-week gestation baby, my heart would want me to do as much as they can to save that baby. But my head says, no, it's not meant to be. And I wouldn't like that child to suffer. And the outcome is quite poor. So I wouldn't want them to pursue any active treatment...

The nurses in this study were all in agreement that it would have been unbearable for them to deliver an extremely premature baby. They held that this was not about not wanting a child with a disability. Rather, it mirrors McHaffie and Fowlie's⁴⁶ finding that nurses did not reject imperfection, they rejected pain and suffering for their own baby. The pain and suffering of extremely premature babies and the effect on neonatal nurses have been explored by Green et al.¹⁹ The nurses in this study were not indifferent to the needs of disabled babies and their families. There is much in the data that reflect the opposite, and for the nurses, if society was going to save these babies then a range of resources and supports needed to be available in the community to assist and enable the parents, and help the child achieve its potential, whatever that potential might be.

Neonatal nurses are the main caregivers in the NICU and they spend the most time with the baby and family. After meeting with the neonatologist, the parents might have further questions when they return to the baby's bedside, and the nurses answer questions and interpret information.⁵³ Gallagher et al.⁵⁴ wondered if neonatal nurses may be a major factor in determining how parents are engaged in decision-making and were not sure if the attitudes that the nurses held in relation to extremely premature babies impacted on their provision of nursing care. This is concerning because conveying unfavourable attitudes and beliefs has the potential to impact decision-making, and that is not within the scope of the role of the neonatal nurse. In fact, there is no evidence in the current research that the nurses even spoke to the parents about their attitudes and beliefs about babies of extreme prematurity. The nurses in this study were adamant that they would never tell the parents what they would or would not do themselves, and as one nurse stated 'I might not want one [extreme prem] myself, but I would never tell a parent' (Nurse 8). Another nurse said 'parents ask me what I would do; I tell them I don't know – of course I know. Well, I think I know' (Nurse 12). Yes, the nurses in this study experienced difficulty providing care at times (Green et al.);^{15,19} however, they went to great pains to emphasise that it did not affect how they cared for the baby or the parents. There was however evidence that the nurses advocated for the baby to ensure it had adequate pain relief (Green et al.).¹⁹

'Imagined futures' has been described by France et al.⁵⁵ who looked at men and women who had terminated a pregnancy because of a foetal abnormality. Experiential knowledge of disability and using

their imagination about the future of their unborn child helped them make their decision.⁵⁵ The use of hypothetical scenarios to investigate the attitude of people (with experiential knowledge of disability, or experience with a disability or contact with a disabled person) to the termination of pregnancy has been shown to have both positive and negative results.⁵⁵ Other studies have shown that expectant parents, who had the antenatal diagnosis of potentially disabling conditions, sought out other people's experiences of disability to help make their decision.⁵⁶⁻⁵⁸ Others were inclined to terminate the pregnancy because of concerns about the child's future and how the family would cope.⁵⁹ Very few studies have examined the decision that people actually take when a pregnancy is affected by a foetal abnormality. France et al.⁵⁵ described a tension between the parent's view of the child's perceived quality of life and the couple's willingness to parent a disabled baby. Although there is no evidence in the data, the authors of this article wonder if this is the reason the nurses would not want to save their baby if they delivered an extremely premature baby. What they have seen in the past and the struggles of parents have firmly planted ideas of what they want or do not want in their own mind.

All of the nurses in this study were very experienced with providing care to the extremely premature baby. They were very knowledgeable about all aspects of care, and they have all seen babies and their parents return to the nursery, and they have seen first-hand what the baby looked like and heard stories from the parents and even the grandparents about how they were managing. It would be interesting to uncover if gender is an issue in this type of hypothetical decision-making. All of the nurses were females, so perhaps a complete picture would only be gained with a male neonatal nursing perspective. Male neonatal nurses were not excluded from the study; a male neonatal nurse did not volunteer to be interviewed. While much has been made of the differences in moral development between males⁶⁰ and females,⁶¹ a meta-analysis of gender differences in moral reasoning by Jaffee and Shibley Hyde⁶² showed only a small difference between males and females in relation to care and justice orientation, meaning that male neonatal nurses might be in agreement with their female colleagues in relation to the hypothetical question under investigation.

In relation to the long-term care of surviving extremely premature babies, extrapolation to the literature shows that females are generally the ones with child rearing responsibilities. It was the understanding of the nurses in this study that it was the mother who provided the majority of care for the baby when discharged. Women provide care in over 75% of situations.⁶³ Children with profound impairment require complex and specialised around-the-clock care, and it is usually, according to Brett⁶⁴ the mother who meets the child's needs. This caregiving could be responsible for the adverse employment outcomes noted in mothers of a child with a disability.⁶⁵ Women caregivers also suffer disproportionately with poorer psychological and physical health.⁶⁶ Could the issue be about subconsciously not burdening the mother or another woman?

The nurses in this study have seen the outcomes or survivors of extreme prematurity. McHaffie and Fowlie⁴⁶ suggest that the NICU is an artificial world and that staff need to glimpse the realities of NICU survivors to increase their understanding of what families endure. The nurses in this study stated that medical and nursing staff were occasionally wrong in their predictions of outcomes, but mostly their predictions were correct. All nurses spoke of observing one baby who had defied predictions and do well, but very few spoke of more than one. Doctors and nurses have been found to be wrong in one-third of their predictions about the baby's survival; however, nearly 90% of those predicted to die but who lived developed severe neurological problems. Doctors and nurses were accurate in their predictions of neurologically intact survival.⁶⁷

In the case of the nurses in the current research, one cannot investigate their real-life decisions, and one can only extrapolate to what they believe they would do in a particular life situation. The conclusions from a hypothetical situation may not always be totally correct, but they could serve as a starting point for further research. Therefore, because a hypothetical is just that there is no reason to preclude a carefully formulated hypothetical thesis if it helps to arrive at a greater understanding. Qualitative method has a valid place in this

research, inasmuch as if researchers want to hear the views of ‘research subjects’ they need to ask them, and give them time to answer fully to explain how they arrive at their understandings.

Hypothetical scenarios are valid in that they are based on hard facts or difficult realities as perceived by the nurses. The ‘facts’ for the nurses regarding neonatal outcomes can change with education and accurate awareness of the latest research and outcome data. Neonatal nurses need to keep updating their education about the survival, morbidity and long-term outcomes of extremely premature babies because neonatal nurses in a study by Blanco et al.¹⁰ underestimated survival rates and overestimated disability rates. The nurses in this study thought long and hard about the outcomes of extremely premature babies and their families, and at times they were distressed by seeing what parents were expected to endure. In order for them to continue to be productive members of the NICU team, the nurses could benefit from clinical supervision, reflection and education with members of the multi-disciplinary team,⁶⁸ where they can work through issues related to the outcomes of extreme prematurity.

Conclusion

Knowledge of the outcomes of extreme prematurity and what they have seen in the past has led the neonatal nurses in this qualitative study informed by phenomenology to believe that they would not want their own baby resuscitated and treated if they delivered at 24 weeks’ gestation and less. The international implications are clear; with increasing technology and refined techniques, the numbers of extremely premature babies will not decrease, and until the outcomes are considered better by the nurses, the primary caregivers of the babies, they would not choose to walk in these parents’ footsteps. Choice was important for the nurses, as was fear of the technological imperative or the snowball effect of technology; it being easier to start treatment and harder to stop. The empathic response of the nurses to the situation of the parents and the baby was authentic, making the nurses believe that the resuscitation and treatment of an extremely premature baby was not the path they would choose to take themselves.

Author contributions

J.G., P.D. and A.A. contributed to study design; J.G. contributed to data collection; J.G., P.D. and A.A. contributed to analysis; and J.G., D.J. and P.D. contributed to manuscript presentation.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship and/or publication of this article.

References

1. Tomlinson MW, Kaempf JW, Ferguson LA, et al. Caring for the pregnant woman presenting at periviable gestation: acknowledging the ambiguity and uncertainty. *Am J Obstet Gynecol* 2010; 202(6): 529.e1–529.e6.
2. Subramanian KNS. Extremely low birth weight infant, pp. 1–21, <http://emedicine.medscape.com/article/979717-overview> (2012, accessed 26 October 2015).
3. Griswold KJ and Fanaroff JM. An evidence-based overview of prenatal consultation with a focus on infants born at the limits of viability. *Pediatrics* 2010; 125(4): e931–e937.
4. Lantos JD and Meadow W. Infants, medical aspects and issues in the care of. In: Post SG (ed.) *Encyclopedia of bioethics*. 3rd ed. New York: MacMillan, 2004, pp. 1252–1257.

5. Moore T, Hennessy EM, Myles J, et al. Neurological and developmental outcome in extremely preterm children born in England in 1995 and 2006: the EPICure studies. *Brit Med J* 2012; 345: e7961.
6. Iacovidou N, Varsami M and Syggellou A. Neonatal outcome of preterm delivery. *Ann NY Acad Sci* 2010; 1205: 130–134.
7. Moster D, Lie RT and Markestad T. Long-term medical and social consequences of preterm birth. *N Engl J Med* 2008; 359: 262–273.
8. Neubauer AP, Voss W and Kattner E. Outcome of extremely low birth weight survivors at school age: the influence of perinatal parameters on neurodevelopment. *Eur J Pediatr* 2008; 167(1): 87–95.
9. Cole CR, Hansen NI, Higgins RD, et al. Very low birth weight preterm infants with surgical short bowel syndrome: incidence, morbidity and mortality, and growth outcomes at 18 to 22 months. *Pediatrics* 2008; 122: e573–e582.
10. Blanco F, Suresh G, Howard D, et al. Ensuring accurate knowledge of prematurity outcomes for prenatal counselling. *Pediatrics* 2005; 115(4): e478–e487.
11. Doyle LW and Anderson PJ. Adult outcome of extremely preterm infants. *Pediatrics* 2010; 126(2): 342–351.
12. Merriam-Webster Dictionary. Hypothetical, <http://www.merriam-webster.com/dictionary/hypothetical> (accessed 14 April 2015).
13. Persky S, Kaphingst KA, Condit CM, et al. Assessing hypothetical scenario methodology in genetic susceptibility testing analog studies: a quantitative review. *Genet Med* 2007; 9(11): 727–738.
14. Green JA. Balancing hope with reality: caregiving dilemmas for neonatal nurses in caring for extremely premature babies. PhD Thesis, Flinders University, Adelaide, SA, Australia, 2007.
15. Green J, Darbyshire P, Adams A, et al. Looking like a proper baby: nurses' experiences of caring for extremely premature infants. *J Clin Nurs* 2015; 24(1–2): 81–89.
16. Green J, Darbyshire P, Adams A, et al. A burden of knowledge: a qualitative study of experiences of neonatal intensive care nurses' concerns when keeping information from parents. *J Child Health Care* 2015; 19(4): 485–494.
17. Green J. Living in hope and desperate for a miracle: NICU nurses perceptions of parental anguish. *J Relig Health* 2015; 54(2): 731–744.
18. Green J, Darbyshire P, Adams A, et al. Balancing hope with reality: how neonatal nurses manage the uncertainty of caring for extremely premature babies. *J Clin Nurs* 2015; 24(17–18): 2410–2418.
19. Green J, Darbyshire P, Adams A, et al. It's agony for us as well Neonatal nurses reflect on iatrogenic pain. *Nurs Ethics* 2016; 23(2): 176–190.
20. Green J, Darbyshire P, Adams A, et al. The myth of the miracle baby: how neonatal nurses interpret media accounts of babies of extreme prematurity. *Nurs Inquiry* 2015; 22(3): 273–281.
21. Green J, Darbyshire P, Adams A, et al. Desperately seeking parenthood: neonatal nurses reflect on parental anguish. *J Clin Nurs* 2015; 24(13–14): 1885–1894.
22. Green J, Darbyshire P, Adams A, et al. Quality versus quantity the complexities of quality of life determinations for neonatal nurses. *Nurs Ethics* 2016. DOI: 10.1177/0969733014558968.
23. Armentrout D. Attitudes/beliefs/feelings held by neonatal nurses toward the care and management of fetal-infants. *Neonatal Netw* 1986; 5(1): 23–29.
24. Van Manen M. *Researching lived experience: human science for an action sensitive pedagogy*. New York: State University of New York Press, 1990, p. 9.
25. Braun V and Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; 3: 77–101.
26. Koch T. Establishing rigour in qualitative research: the decision trail. *J Adv Nurs* 2006; 53: 91–100.
27. Stinson R and Stinson P. *The long dying of baby Andrew*. Boston, MA: Little, Brown and Co., 1983.
28. Mayo Clinic. Premature birth, <http://www.mayoclinic.org/diseases-conditions/premature-birth/basics/symptoms/con-20020050> (accessed 14 June 2105).
29. Smith L, Draper ES and Field D. Long-term outcome for the tiniest or most immature babies: survival rates. *Semin Fetal Neonatal Med* 2014; 19(2): 72–77.

30. Saigal S and Doyle LW. An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet* 2008; 371(9608): 261–269.
31. Blencowe H, Cousens S, Oestergaard MZ, et al. National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet* 2012; 379(9832): 2162–2172.
32. Government of South Australia. *South Australian perinatal practice guidelines: perinatal care at the threshold of viability*. Adelaide, SA, Australia: South Australian Maternal and Neonatal Clinical Network, 2013, pp. 1–14.
33. Queensland Government. *Queensland clinical guideline: perinatal care at the threshold of viability*, 2014, pp. 1–35, <https://www.health.qld.gov.au/qcg/documents/g-viability.pdf> (accessed 14 August 2016).
34. Larcher V. Ethical considerations in neonatal end-of-life care. *Semin Fetal Neonatal Med* 2013; 18(2): 105–110.
35. Weiss AR, Binns HJ and Collins JW. Decision-making in the delivery room: a survey of neonatologists. *J Perinatol* 2007; 27(12): 754–760.
36. Donohue PK, Boss RD, Aucott SW, et al. The impact of neonatologists' religiosity and spirituality on health care delivery for high-risk neonates. *J Palliat Med* 2010; 13(10): 1219–1224.
37. Lawrence RE and Curlin FA. Autonomy, religion and clinical decisions: findings from a national physician survey. *J Med Ethics* 2009; 35(4): 214–218.
38. Paris JJ, Schreiber MD and Elias-Jones A. Resuscitation of the preterm infant against parental wishes. *Arch Dis Child Fetal Neonatal Ed* 2005; 90(3): F208–F210.
39. Kühnberger A, Schulte-Mecklenbeck M and Perner J. Framing decisions: hypothetical and real. *Organ Behav Hum Dec* 2002; 89(2): 1162–1175.
40. D'Argembeau A and Mathy A. Tracking the construction of episodic future thoughts. *J Exp Psychol Gen* 2011; 140(2): 258.
41. Sanders MR, Donohue PK, Oberdorf MA, et al. Perceptions of the limit of viability: neonatologists' attitudes toward extremely preterm infants. *J Perinatol* 1994; 15(6): 494–502.
42. Streiner DL, Saigal S, Burrows E, et al. Attitudes of parents and health care professionals toward active treatment of extremely premature infants. *Pediatrics* 2001; 108(1): 152–157.
43. De Leeuw R, Cuttini M, Nadai M, et al. Treatment choices for extremely preterm infants: an international perspective. *J Pediatr* 2000; 137(5): 608–616.
44. Oei J, Askie LM, Tobiensky R, et al. Attitudes of neonatal clinicians towards resuscitation of the extremely premature infant: an exploratory survey. *J Paediatr Child Health* 2000; 36(4): 357–362.
45. Janvier A, Leblanc I and Barrington KJ. Nobody likes premies: the relative value of patients' lives. *J Perinatol* 2008; 28(12): 821–826.
46. McHaffie HE and Fowlie PW. *Life, death and decisions: doctors and nurses reflect on neonatal practice*. Cheshire: Hochland & Hochland Ltd, 1996.
47. Janvier A, Lantos J, Deschenes M, et al. Caregivers attitudes for very premature infants: what if they knew? *Acta Paediatr* 2008; 97(3): 276–279.
48. Janvier A, Nadeau S, Deschenes M, et al. Moral distress in the neonatal intensive care unit: caregiver's experience. *J Perinatol* 2007; 27(4): 203–208.
49. Lantos JD, Miles SH, Silverstein MD, et al. Survival after cardiopulmonary resuscitation in babies of very low birth weight: is CPR futile therapy? *N Engl J Med* 1988; 318(2): 91–95.
50. Barr P and Courtman SP. Cardiopulmonary resuscitation in the newborn intensive care unit. *J Paediatr Child Health* 1998; 34(6): 503–507.
51. McHaffie HE and Fowlie PW. Should the baby live? The role of nurses in decision making. *NT Research* 1997; 2(4): 271–279.
52. Harrison H. Neonatal intensive care: parent's role in ethical decision making. *Birth* 1986; 13(3): 165–175.

53. Douglas SM and Dahnke MD. [Creating an ethical environment for parents and health providers dealing with the treatment dilemmas of neonates at the edge of viability.](#) *J Neonatal Nurs* 2013; 19(1): 33–37.
54. Gallagher K, Marlow N, Edgley A, et al. [The attitudes of neonatal nurses towards extremely preterm infants.](#) *J Adv Nurs* 2012; 68(8): 1768–1779.
55. France EF, Locock L, Hunt K, et al. [Imagined futures: how experiential knowledge of disability affects parents' decision making about fetal abnormality.](#) *Health Expect* 2012; 15(2): 139–156.
56. Chaplin J, Schweitzer R and Perkoulidis S. [Experiences of prenatal diagnosis of spina bifida or hydrocephalus in parents who decide to continue with their pregnancy.](#) *J Genet Couns* 2005; 14(2): 151–162.
57. Redlinger-Grosse K, Bernhardt BA, Berg K, et al. [The decision to continue: the experiences and needs of parents who receive a prenatal diagnosis of holoprosencephaly.](#) *Am J Med Genet* 2002; 112(4): 369–378.
58. Petrucelli N, Walker M and Schorry E. [Continuation of pregnancy following the diagnosis of a fetal sex chromosome abnormality: a study of parents' counseling needs and experiences.](#) *J Genet Couns* 1998; 7(5): 401–415.
59. Sawyer SM, Cerritelli B, Carter LS, et al. [Changing their minds with time: a comparison of hypothetical and actual reproductive behaviors in parents of children with cystic fibrosis.](#) *Pediatrics* 2006; 118(3): e649–e656.
60. Kohlberg L. Stage and sequence. In: Goslin DD (ed.) *Handbook of socialization theory and research*. Chicago, IL: Rand McNally, 1969, pp. 348–480.
61. Gilligan C. *In a different voice*. Cambridge, MA: Harvard University Press, 1982.
62. Jaffee S and Hyde JS. [Gender differences in moral orientation: a meta-analysis.](#) *Psychol Bull* 2000; 126(5): 703–726.
63. Kasuya RT, Polgar-Bailey MP and Robbyn Takeuchi MSW. [Caregiver burden and burnout. A guide for primary care physicians.](#) *Postgrad Med* 2000; 108(7): 119.
64. Brett J. [The experience of disability from the perspective of parents of children with profound impairment: is it time for an alternative model of disability?](#) *Disabil Soc* 2002; 17(7): 825–843.
65. Baker DL and Drapela LA. [Mostly the mother: concentration of adverse employment effects on mothers of children with autism.](#) *Soc Sci J* 2010; 47(3): 578–592.
66. Pinquart M and Sörensen S. [Gender differences in caregiver stressors, social resources, and health: an updated meta-analysis.](#) *J Gerontol B Psychol Sci Soc Sci* 2006; 61(1): P33–P45.
67. Frain L, Ren Y, Meadow A, et al. [Certainty, uncertainty, and accuracy of daily predictions of non-survival in the NICU: we're often wrong but rarely in doubt.](#) *Pediatr Res* 1998; 43: 29–39.
68. Brunero S and Stein-Parbury J. [The effectiveness of clinical supervision in nursing: an evidenced based literature review.](#) *Aust J Adv Nurs* 2008; 25(3): 86–94.