

Increased survival means increasing roles for primary care after cancer diagnosis

Rosalind Adam, Niek de Wit, Patti Groome, Charles Helsper, Mary McBride, Eila Watson and Jan Wind

Morgan et al point out that cancer survival rates in the UK are improving.¹ It is timely to consider the roles that GPs play following a cancer diagnosis. We would like to report highlights from a workshop (Cancer in Primary Care International Network [Ca-PCI] annual scientific meeting, Edinburgh, 2017), which explored the roles of GPs following a cancer diagnosis in the Netherlands, Canada, the UK, and Australia, and showcased current improvement initiatives and research.

Internationally, the roles that GPs play in cancer care are expanding,² but GP input during treatment and survivorship phases is highly variable within and between countries.³ In the UK, a formal cancer care review is remunerated under the current GP contract, but there is evidence that the review is often unstructured, and is perceived to be of limited use.⁴ Lack of standardised approaches in primary care following a cancer diagnosis are problematic not only because of the dramatic increase in cancer prevalence, but also because of the increasingly chronic nature of the disease and the high prevalence of comorbid diseases. Consequently, more cancer patients and their partners consult their GPs more frequently over longer periods of time.⁵ There is a political and professional will to provide comprehensive, cost-effective care following a cancer diagnosis, and a sense that primary care is uniquely placed to contribute to this.

Indicators include improving patient–professional communication, shared decision making, and continuity of primary care after a cancer diagnosis;⁶ developing guidelines to standardise survivorship care (and promoting adherence to existing guidelines); developing specific primary-care-based behavioural and lifestyle interventions to improve outcomes in cancer survivors; improving primary/secondary care communication; and developing digital systems to support clinical information exchange, patient self-management, for example, the ‘OncoKompas’,⁷ and to improve recurrence detection, for example, the ASSICA melanoma intervention.⁸

REFERENCES

1. Morgan I, Wilkes S (2017) Br J Gen Pract, Improving early diagnosis of cancer in UK general practice. DOI: <https://doi.org/10.3399/bjgp17X691265>.Google Scholar
2. Rubin G, Berendsen A, Crawford SM, et al. (2015) The expanding role of primary care in cancer control. Lancet Oncol 16(12):1231–1272.Google Scholar
3. Jiang L, Lofters A, Moineddin R, et al. (2016) Primary care physician use across the breast cancer care continuum: CanIMPACT study using Canadian administrative data. Can Fam Physician 62(10):e589–e598.Google Scholar
4. Walter FM, Usher-Smith JA, Yadlapalli S, Watson E (2015) Caring for people living with, and beyond, cancer: an online survey of GPs in England. Br J Gen Pract doi:10.3399/bjgp15X687409.Google Scholar

5. Heins M, Schellevis F, Rijken M, et al. (2013) Partners of cancer patients consult their GPs significantly more often with both somatic and psychosocial problems. *Scand J Prim Health Care* 31(4):203–208. [Google Scholar](#)
6. de Wit NJ (2017) A ‘time out consultation’ in primary care for elderly patients with cancer: better treatment decisions by structural involvement of the general practitioner. *Eur J Cancer Care* doi:10.1111/ecc.12711. [Google Scholar](#)
7. Duineveld LA, Wieldraaijer T, van Asselt KM, et al. (2015) Improving care after colon cancer treatment in the Netherlands, personalised care to enhance quality of life (I CARE study): study protocol for a randomised controlled trial. *Trials* 16:284. [Google Scholar](#)
8. Murchie P, Allan JL, Brant W, et al. (2015) Total skin self-examination at home for people treated for cutaneous melanoma: development and pilot of a digital intervention. *BMJ Open* 5(8):e007993.