

***De novo* variants in *MED12* cause X-linked syndromic neurodevelopmental disorders in 18 females**

D.L. Polla^{1,2}, E.J. Bhoj PhD³, J.B.G.M. Verheij MD⁴, J.S. Klein Wassink-Ruiter MD⁴, A. Reis MD⁵, C. Deshpande⁶, A. Gregor PhD⁵, K. Hill-Karfe³, A.T. Vulto-van Silfhout MD PhD⁷, R. Pfundt PhD⁷, E.M.H.F. Bongers MD PhD⁷, H. Hakonarson MD PhD³, S. Berland MD⁸, G. Gradek MD⁸, S. Banka MD PhD^{9,10}, K. Chandler⁹, L. Gompertz⁹, S.C. Huffels¹, C.T.R.M. Stumpel MD¹¹, R. Wennekes¹², A.P.A. Stegmann PhD¹¹, W. Reardon MD¹³, E.K.S.M. Leenders⁷, B.B.A. de Vries MD PhD⁷, D. Li PhD¹⁴, E. Zackai MD³, N. Ragge MD¹⁵, S.A. Lynch¹⁶, S. Cuddapah MD³, H. van Bokhoven PhD¹, C. Zweier MD PhD^{5*}, A.P.M. de Brouwer PhD^{1*}.

¹ Department of Human Genetics, Donders Institute for Brain, Cognition and Behavior, Radboud University Medical Center, Nijmegen, The Netherlands.

² CAPES Foundation, Ministry of Education of Brazil, Brasília, Brazil.

³ Department of Pediatrics, Division of Human Genetics and Molecular Biology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA.

⁴ Department of Medical Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

⁵ Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany.

⁶ Department of Clinical Genetics, SE Thames Regional Genetics Service, Guy's Hospital, London, United Kingdom.

⁷ Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands.

⁸ Department of Medical Genetics, Haukeland University Hospital, Helse Bergen, Norway

⁹ Manchester Centre For Genomic Medicine, University of Manchester, St Mary's Hospital, Manchester Academic Health Science Centre, Manchester, UK.

¹⁰ Division of Evolution & Genomic Sciences, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester M13 9PL, UK.

¹¹ Department of Clinical Genetics and GROW-School for Oncology and Developmental Biology, Maastricht University Medical Center, Maastricht, The Netherlands.

¹² Department of Radiology, Maastricht University Hospital, Maastricht University, Maastricht, The Netherlands.

¹³ Department of Clinical Genetics. Children's Health Ireland, Dublin, Ireland.

¹⁴ Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, PA, USA

¹⁵ International Centre for Eye Health, London School of Hygiene and Tropical Medicine, London, United Kingdom.

¹⁶ UCD Academic Centre on Rare Diseases, School of Medicine and Medical Sciences, University College Dublin, and Clinical Genetics, Temple Street Children's University Hospital, Dublin, Ireland.

* , these authors contributed equally

Correspondence to: Arjan.deBrouwer@radboudumc.nl

Abstract

Purpose: MED12 is a subunit of the Mediator multiprotein complex with a central role in RNA polymerase II transcription and regulation of cell growth, development, and differentiation.

This might underlie the variable phenotypes in males carrying missense variants in *MED12*, including X-linked recessive Ohdo-, Lujan-, and FG syndromes.

Methods: By international matchmaking we assembled variant and clinical data on 18 females presenting with variable neurodevelopmental disorders (NDDs) and harboring *de novo* variants in *MED12*.

Results: Five nonsense variants clustered in the C-terminal region, two splice variants were found in the same exon 8 splice acceptor site, and 11 missense variants were distributed over the gene/protein. Protein truncating variants were associated with a severe, syndromic phenotype consisting of intellectual disability (ID), facial dysmorphism, short stature, skeletal abnormalities, feeding difficulties and variable other abnormalities. *De novo* missense variants were associated with a less specific, but homogeneous phenotype including severe ID, autistic features, limited speech and variable other anomalies, overlapping both with females with truncating variants as well as males with missense variants.

Conclusion: We establish *de novo* truncating variants in *MED12* as causative for a distinct NDD and *de novo* missense variants as causative for a severe, less specific NDD in females.

Keywords: MED12, intellectual disability, neurodevelopmental disorders, X-linked, female.

Introduction

MED12 is one of 31 subunits (MED1-MED31) of the large multiprotein Mediator complex that regulates gene expression in all eukaryotes by interacting with RNA polymerase II.^{1,2} It is involved in transcriptional elongation and termination, mRNA processing, noncoding RNA activation, super enhancer formation, and epigenetic regulation.^{2,3} Hence, Mediator complex is a master coordinator of cell growth and homeostasis, development, and differentiation,⁴ and its subunit MED12 is a critical transducer of regulatory information essential for organogenesis.⁵ MED12 consists of the Med12 domain that forms part of the kinase section of Mediator, which can negatively regulate the Gli3-dependent sonic hedgehog signaling pathway via its interaction with Gli3 within RNA polymerase II,⁶ the Med12-LCEWAV domain, of which the function is not known, and the Med12-PQL domain that binds to β -catenin, which targets MED12 to activate transcription of Wnt-responsive genes.⁷

Hemizygous missense variants in *MED12* are known to cause at least three different but overlapping X-linked syndromic neurodevelopmental disorders (NDDs) in males: FG syndrome (FGS1, also known as Opitz-Kaveggia syndrome [OKS; OMIM 305450]), Lujan-Fryns syndrome (LS; OMIM 309520), and X-linked Ohdo syndrome (OHDOX; OMIM 300895). FGS1 is characterized by intellectual disability (ID), hypotonia, dysmorphic features such as macrocephaly, prominent forehead, downslanting palpebral fissures, small ears with occasionally sensorineural hearing loss, broad thumbs and halluces, constipation and/or anal anomalies, partial agenesis of the corpus callosum and characteristic hyperactive and talkative behavior.^{8,9} LS is characterized by mild to moderate ID, dysmorphic features such as macrocephaly, large head, a long thin face, low-set ears, prominent nasal bridge, high narrow palate, short and deep philtrum, tall stature with thin or marfanoid habitus, behavioral aspects and abnormalities of the corpus callosum.^{10,11} OHDOX is characterized by ID and typical facial features, including blepharophimosis, facial coarsening, thick alae nasi, and a triangular face.^{12,13} Additionally, variants in *MED12* have been implicated in non-specific NDDs in males.^{11,14–28} Though carrier females in these families are usually not affected, three

females with inherited *MED12* variants,^{15,17,25} p.(Ser1967Glnfs*84), p.(Arg1148His) and p.(Ile771Thr), and three females with *de novo* *MED12* variants,^{29–31} p.(Glu172Gln), c.1249-1G>C and p.(Arg521His), were reported to have NDDs with variable cognitive impairment.

Here, we report on assembled clinical and genetic data of 18 females with *de novo* variants in *MED12*. Truncating variants in seven individuals resulted in a syndromic presentation with variable ID, short stature, facial dysmorphism, feeding difficulties and skeletal abnormalities. Missense variants in 11 individuals were associated with a phenotype of severe ID, autistic features and limited speech and variable dysmorphic features.

Materials and Methods

DNA analysis and X-inactivation

Affected individuals were referred by physicians working in disability services and clinical geneticists, and connected through personal communication, Decipher database³² and GeneMatcher.³³ *MED12* variants were identified by exome or genome sequencing and confirmed by Sanger sequencing (**Supplementary Table 1**). X-chromosome inactivation (XCI) analysis was performed at the androgen receptor locus with a modification of the assay previously described.³⁴ Primer sequences and PCR conditions for Sanger sequencing and XCI testing are available upon request.

RNA Isolation, cDNA Synthesis and RT-PCR

All RNA isolations were performed with the NucleoSpin RNA Clean-up Kit (cat. no. 740955-50, Macherey-Nagel, Düren, Germany) according to manufacturer's protocol. RNA was quantified by nanodrop. One microgram of total RNA was used for all cDNA synthesis reactions. For reverse transcription-PCR (RT-PCR) analysis, cDNA was synthesized by the SuperScript VILO Master Mix (cat. no. 11755050, Thermo Fisher Scientific, Waltham, MA, USA), following manufacturer's instructions.

To determine the potential splice acceptor variant effect on *MED12* splicing in individuals 6 and 7, RT-PCR analysis was performed using primers located in exon 8 (forward) and exon 12 (reverse) of *MED12*. Primer sequences are available upon request. For RT-PCR analysis, 25 ng of cDNA was used for all reactions. Amplifications were performed according to standard procedures. PCR products were sequenced using the ABI PRISM BigDye Terminator Cycle Sequencing V2.0 Ready Reaction Kit and analyzed with the ABI PRISM 3730 DNA analyzer (Applied Biosystems, Foster City, CA, USA).

Minigene-based splice assay

The effect of the two splice acceptor variants was also assessed by minigene-based splicing assays employing wild-type and mutant constructs using DNA from individuals 6 and 7 as a template. To amplify the genomic region encompassing exons 8 and 9 of the *MED12* gene, we designed PCR primers located in exons 7 (forward) and 11 (reverse) using Primer3plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi>). The PCR reaction mixtures of 50 µl contained 25 µM of each primer pair, a high-fidelity Taq polymerase Phusion® High-Fidelity DNA Polymerase (cat. no. M0530L, New England BioLabs, Ipswich, MA, USA), 5X PCR HF buffer, 1µl of dNTP mix, 5X Q solution and 5ng genomic DNA from patients 1 and 2. Amplifications were performed using a two-phase touchdown PCR program described before. The resulting PCR products were run on a 0.8% agarose gel and were purified with the NucleoSpin Gel and PCR cleanup kit (cat. no. 740609.250; Macherey-Nagel, Düren, Germany), according to the manufacturer's protocol. One hundred fifty nanograms of each purified insert were cloned into the pDONR201 vector by using the Gateway BP Clonase Enzyme Mix (cat. no. 11789021; Thermo Fisher Scientific, Waltham, MA, USA).

The different pDONR cloned constructs were validated by Sanger sequencing, and 150 ng of each were cloned into the destination vector pCI-NEO-RHO exon3,5/DEST by using the Gateway LR Clonase enzyme mix (cat. no. 11791043; Thermo Fisher Scientific, Waltham, MA, USA). Three minigene constructs were generated, one with the wild-type and

two with the c.1249-1G>C and c.1249-2A>G variants. HEK293T cells (ATCC CRL-3216; American Type Culture Collection [ATCC], Wesel, Germany) were grown in DMEM supplemented with 10% (volume/volume) fetal calf serum (cat. no. F7524; Sigma Aldrich, St. Louis, MO, United States), 1% (volume/volume) Sodium pyruvate (cat. no. S8636; Sigma Aldrich, St. Louis, MO, United States), and penicillin/streptomycin (cat. no. P4333; Sigma Aldrich, St. Louis, MO, United States). Cells were seeded in 6-well plates at a density of 5×10^5 cells/well and transfected the day after with 2 μ g of each minigene using FuGENE HD Transfection Reagent (cat. no. E2311; Promega, Madison, WI, USA) with a DNA-to-liposome ratio of 1 μ g:3 μ l, according to the manufacturer's instructions. Medium was replaced after 24 h, and cells were harvested 48 h after transfection. Total RNA isolation, cDNA production and RT-PCR reactions were performed as described before. The RT-PCR products were resolved on a 2% (weight/volume) agarose gel, the resulting bands were excised, and the nucleic acid was purified by using NucleoSpin Gel & PCR cleanup (cat. no. 740609.50, Macherey-Nagel, Düren, Germany). Sanger sequencing was performed on 100 ng of each purified band by using the same primers used in the PCR reaction showing no other variant except those introduced by us.

Results

Variant spectrum, splicing effects and X-inactivation

Five individuals carried *de novo* nonsense variants in the last four exons and clustering in or down-stream of the catenin binding domain of *MED12* (NM_005120.2) (**Fig. 1**). In two individuals, *de novo* variants affecting the splice acceptor of exon 8 were identified: c.1249-1G>C and c.1249-2A>G. Additionally, 11 females carried *de novo* missense variants predicted to be deleterious (**Supplementary Table 1**). The identified missense variants were distributed over the N-terminal two thirds of the protein, and all but one were located between the LCEWAV and the catenin binding domain (**Fig. 1**). p.(Arg1138Trp) recurrently occurred in two individuals. None of the identified variants were present in the Genome Aggregation Database (gnomAD; <https://gnomad.broadinstitute.org>; **Supplementary Table 1**). The *MED12* pLi score was 1 and the Z-score was 6.58, which indicates that there is strong selection against protein truncating variants and missense variants in males and females.

We used RT-PCR to investigate whether the two variants in the canonical splice acceptor of exon 8 affected splicing. We found expression of only the wild-type fragment in EBV-LCL and fibroblast lines from individual 1 carrying the c.1249-1G>C variant, and in a fibroblast line from individual 2 who carried the c.1249-2A>G variant (**Fig. 2B-C**). To investigate the splicing defect in more detail, we generated two *MED12* minigenes spanning exons 7 to 11 using control and patient genomic DNAs. HEK293T cells were transfected with mutant and control minigenes (**Fig. 2D**). While the control minigene did not show any splicing defect, both mutated minigenes showed complete skipping of exon 9 (**Fig. 2E**), leading to an abnormal transcript due to an early termination codon: p.(Val417Alafs*76). This indicates that the presence of only the wildtype allele in fibroblasts and EBV-LCLs might be due to either nonsense-mediated mRNA-decay or more likely skewing of XCI.

Analysis of the X inactivation status showed extreme skewing (>95:5) in blood or fibroblasts of three individuals with protein truncating variants including individual 2, and random XCI in three individuals (**Supplementary Table 1**). In individual 1, the two androgen

receptor alleles were non-informative. In nine out of the 11 individuals with missense variants, XCI was extremely skewed (>95%) and skewed (85%) in one. In three individuals with a missense variant, XCI was random in peripheral blood cells. In four individuals where two different tissues, whole blood and either a fibroblast cell line or EBV-LCLs, were tested, XCI behaved comparable between tissues. Three of these individuals had skewing of X-inactivation, and the variant was located on the inactive allele in the tested tissues.

Clinical spectrum

All seven individuals with *de novo* protein truncating variants in *MED12* presented with ID, ranging from mild in one to severe/profound in four individuals (**Table 1**; **Supplementary Table 1**). Speech was commonly delayed, and absent in two individuals. Brain imaging by MRI or ultrasound was performed in five individuals and showed partial hypoplasia or agenesis of the corpus callosum and/or enlarged ventricles in three of them. Feeding difficulties were reported in five individuals, muscular hypotonia in four, autistic features in three, and seizures in only one individual. Birth weight and/or length below -2 SD were observed in two individuals as well as congenital microcephaly. Postnatally, short stature occurred in four of six individuals, of whom these data were available. The two individuals who were of normal length were the youngest of the six. This suggests that short stature is a progressive feature that might become apparent only at a later age. Microcephaly and macrocephaly were each reported once, whereas three individuals had a normal head circumference. Skeletal abnormalities were present in six out of the seven individuals and included syndactyly of fingers II-IV or of toes II and III as well as other finger anomalies, scoliosis or rhizomelic limb shortening. Facial gestalt was quite distinct in all the individuals and included sparse hair, low-set, posteriorly rotated ears, a prominent forehead or bitemporal narrowing, hypertelorism, downslanting palpebral fissures, blepharophimosis, palatal anomalies or a bifid uvula, dental crowding, and micro- or retrognathia (**Fig. 3**). Variable other features were laryngeal anomalies requiring tracheostoma in two individuals, anteriorly placed anus (4/7), hearing loss (2/3) and pigmentation anomalies (2/3). Individuals

1 and 2 with variants in the splice acceptor of exon 8 had less prominent micro/retrognathia and shared other clinical aspects such as a very prominent forehead with large fontanelles, hypertelorism and MRI anomalies including partial agenesis of corpus callosum, enlarged ventricles and abnormal tectum and cerebellar vermis.

In contrast, the individuals with *de novo* missense variants in *MED12* presented with a less distinct, but still homogenous phenotype. Eight out of 11 individuals presented with severe to profound ID and six of those with absent speech. One exception was individual 14 with a low-normal IQ of 83, but she had other hallmarks of *MED12*-related disorders, such as thin corpus callosum, a prominent forehead and ptosis. Behavioral abnormalities such as autistic features or attention deficit were common and reported in eight individuals. Corpus callosum anomalies were reported in four and cortical thinning in one of ten individuals in whom MRI was performed. Growth parameters were variable with mostly normal measurements at birth and postnatally. Short stature was reported in only one individual, while tall stature occurred in two individuals. Syndactyly of fingers or toes was reported in two individuals only, while scoliosis and pectus anomalies were reported in three individuals. Ophthalmological anomalies were found in nine individuals of which strabismus and nystagmus were most frequent. Further variable anomalies in single or few individuals included muscular hypotonia, hearing loss or constipation. Subtle and non-specific facial dysmorphisms were noted in several of the individuals (**Fig. 3**).

Discussion

In this study, we identified 18 unrelated females with *de novo* variants in *MED12* and presenting with variable NDDs. These cases can be classified into two molecularly and phenotypically distinct groups. Truncating variants were identified in seven individuals and resided either in the splice acceptor of exon 8 (n=2), resulting in out of frame skipping of exon 9, or clustered C-terminally in or downstream of the catenin binding domain (n=5). The phenotype of these individuals appeared rather distinct and comprised mainly severe ID, short stature, (severe) feeding difficulties, syndactyly of toes II and III and/or syndactyly of fingers II, III and IV and variable other skeletal abnormalities. Marked facial dysmorphism included an abnormal skull shape, sparse hair, low-set, posteriorly rotated ears, blepharophimosis, micro- and/or retrognathia and dental crowding. The presence of abnormal transcripts and subsequently truncated protein might be expected for these seven variants. Particularly for the C-terminal variant p.(Gln2150*), a truncating effect is likely since localization in the last or pen-ultimate exon usually does not lead to nonsense-mediated mRNA decay.³⁵

So far, pathogenic *MED12* variants described in the literature have been mainly missense. Only two other truncating variants were reported previously. Firstly, a frameshift variant, c.5898dupC, also located in the C-terminal catenin-binding domain, was identified in 10 affected males and an affected female from one family with severe to profound ID and facial dysmorphism.¹⁵ Nine carrier females showed only borderline to mild ID. This variant has been demonstrated to escape nonsense-mediated mRNA decay and lead to creation of an additional abnormal transcript due to activation of two cryptic splice sites in exon 41.¹⁵ Secondly, a *de novo* splice variant that is identical to the one identified in individual 1 in this study, was reported in a one year old female with a similar phenotype comprising developmental delay, facial dysmorphism, cleft lip and palate and a thin corpus callosum.³¹ For the two novel splice-variants we identified, we could not detect an aberrant transcript. This could be due to the skewed X-inactivation in the peripheral blood samples of these individuals as we could observe a protein truncating transcript in a minigene assay.

Clustering of the truncating variants in two distinct regions of the gene/protein points to specific effects. The very C-terminal variants leave all known functional domains intact, while the N-terminal splice variants in the splice acceptor of exon 8 would be expected to affect the LCEWAV and the Catenin-binding domain. Individuals 1 and 2 with these splice variants share particular clinical aspects such as skull shape and MRI abnormalities. Apart from that, the phenotype between individuals with C-terminal nonsense variants and N-terminal splice variants is highly overlapping. We suspect a common loss-of-function consequence of the truncating variants rather than a dominant-negative effect. This would be in accordance with gnomAD constraint scores, indicating intolerance of *MED12* towards loss-of-function variants. As truncating variants in *MED12* were mainly observed in females with a severe and syndromic phenotype, their functional consequence might be possibly more severe than the effect of most pathogenic missense variants.

The phenotype in the 11 females with *de novo* missense variants appeared less specific in comparison, but was still homogenous with severe or profound ID in all but two individuals, frequent autistic features and lack or severe limitation of speech. Variable other anomalies included agenesis of corpus callosum, mild short stature, syndactyly, subtle facial dysmorphism and feeding difficulties. Similarly to previously reported missense variants found in males, the *de novo* missense variants identified in females in this study were distributed all over the protein. Previously, only two missense variants, p.(Glu172Gln)²⁹ and p.(Arg521His),³¹ have been reported in females with comparable phenotypes.

Pathogenic variants in *MED12* are associated with various non-syndromic and syndromic NDDs in both males and females. Apart from shared intellectual disability, dysmorphic features such as blepharophimosis, brain anomalies such as agenesis or hypoplasia of corpus callosum or other morphological anomalies such as syndactyly overlap between both males and females but also within the male and within the female NDDs. In addition to the functional consequences (e.g. truncating versus missense variants) and the location of the variants, XCI patterns in females might also contribute to the variable phenotypic expression. Skewing of XCI in favor of the wildtype allele can accordingly

contribute to the compensation of X-chromosomal variants in heterozygous females, thus preventing clinical manifestation. However, XCI patterns in previous reports did not correlate with presence or severity in female carriers of *MED12* variants.^{14,15,17,25,29} While XCI skewing was demonstrated more frequently in 12 individuals with either truncating or missense variants, also random XCI was reported in four individuals, who did not present with a less or more severe phenotype. X-inactivation is a random transcriptional silencing process in gastrulation, where one of the two X chromosomes is inactivated. Afterwards this specific pattern is maintained in the daughter cells that all have the same maternal or paternal X-inactivated, often resulting in functional mosaicism with the mutant allele being active or inactive depending on the tissue. In blood or fibroblast cells of affected individuals, X-inactivation was skewed, and only the wildtype allele could be detected, which would in principle point to a compensatory effect. Other tissues were not available. We therefore speculate that, in contrast to blood, the mutant allele might be active in progenitor brain cells and that this might underly the severe phenotype. Pigmentation anomalies can be suggestive for such a functional mosaicism.³⁶ Such skin pigmentation anomalies were observed in at least three individuals in our cohort, and partial skin depigmentation was reported in one of the previously published female individuals.²⁹

In conclusion, we demonstrate that *de novo* missense and specific truncating variants in *MED12* cause NDDs with variable abnormalities and dysmorphic features in females, partly overlapping with the previously known phenotypic spectrum from affected males. We speculate that “severity”, localization and functional consequences of the variants as well as XCI patterns in females might contribute to the clinical presentation and variability both among females and between affected females and males.

Acknowledgements

We are grateful to the patients and their families for their participation. This work was supported by the European Union's Seventh Framework Program (Gencodys; grant 241995 to HvB). D.L.P. is recipient of a CAPES Fellowship (99999.013311/2013-01). C.Z. was supported by grants from the German Research Foundation (DFG) (ZW184/3-1, ZW184/6-1 and 270949263/GRK2162) and by the IZKF Erlangen (E31). B.B.A.d.V was supported from the Dutch Organization for Health Research and Development (ZON-MW grants 917–86–319 and 912–12–109). C.T.R.M.S. and A.P.A.S. are supported by the European Reference Network (ERN) ITHACA (Project ID no 739543). The 100,000 Genomes Project is funded by the National Institute of Health Research and NHS England. The Wellcome Trust, Cancer Research UK and the Medical Research Council have also funded research structure. This study makes use of data generated by the Genomics England Research Consortium, DDD study and the DECIPHER community. A full list of centres who contributed to the generation of the data is available from <https://decipher.sanger.ac.uk> and via email from decipher@sanger.ac.uk. Funding for the project was provided by Wellcome Trust.

Ethics Declaration

Individuals were identified in different centers worldwide in diagnostic or research settings approved by the respective institutional review boards in each case. The study as a whole was approved by the review board of the Friedrich-Alexander-University Erlangen-Nürnberg. Written informed consent was obtained for all individuals involved. For publishing the patients' photos, specific written informed consent was obtained as well. This study adhered to the World Health Association Declaration of Helsinki (2013).

Conflict of interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Data Availability

All data and materials that is not included in the paper is available upon request.

Author contributions

Conceptualization: A.d.B., A.R. C.Z., H.v.B.; Data Curation: A.d.B, C.Z. D.L.P.; Formal Analysis: D.L.P.; Funding Acquisition: A.P.A.S., B.B.A.d.V, C.T.R.M.S.,C.Z., D.L.P., H.v.B; Investigation: A.G., A.P.A.S., A.T.V., B.B.A.d.V., C.D., C.T.R.M.S., C.Z., D.L., D.L.P., E.J.B., E.K.S.M.L., E.M.H.F.B., E.Z., H.H., G.G., J.B.G.M.V., J.S.K., K.C., K.H-K., L.G., N.R., R.W., S.A.L., S.Ba., S.Be., S.C., S.C.H., W.R. Methodology: R.P. Supervision: A.d.B., A.R., C.Z. H.v.B. Visualization; A.d.B., C.Z., D.L.P. Writing: A.d.B., C.Z., D.L.P. Writing – review & editing: A.d.B., C.Z., D.L.P.

Figure legends

Fig. 1 - Graphical representation of *de novo* and inherited variants distributed across the known functional domains of the MED12 protein.

Known ID-associated missense variants in males are displayed below the protein scheme.

Above are previously reported inherited frameshift,¹⁵ inherited^{17,25} and *de novo*^{29,30} missense, and *de novo* splice site³¹ variants in female individuals, together with the *de novo* variants in *MED12* (NM_005120.2) identified in this study (in bold). MED12 domains; Med12:

Transcription mediator complex subunit Med12. This subunit forms part of the kinase section of Mediator, which can negatively regulate the Gli3-dependent sonic hedgehog signaling pathway via its interaction with Gli3 within the RNA polymerase II. A complex is formed between Med12, Med13, CDK8 and CycC which is responsible for suppression of transcription.⁶ Med12-LCEWAV; Eukaryotic Mediator 12 subunit domain: The function of this region is not known, but there is a conserved sequence motif (LCEWAV). Med12-PQL; Eukaryotic Mediator 12 catenin-binding domain: β -catenin physically and functionally targets this PQL (proline-, glutamine-, leucine-rich) region of the Med12 subunit of Mediator to activate transcription of Wnt-responsive genes.⁷

Fig. 2 – De novo canonical splice site variants in MED12.

(A) Position of the variants (red circle) at the donor site of *MED12* exon 9. Blue arrows represent primers used for analysis. (B, C) No effect on splicing is found on RT-PCR analysis of expression of *MED12* in EBV-LCLs and fibroblast cell lines from patients and controls. One µg cycloheximide (CHX) per ml medium was added for four hours to inhibit nonsense-mediated mRNA decay. (D, E) Minigene analysis shows that the canonical splice site variants induce skipping of exon 9 in the transcript containing the variant. This causes an ORF (open reading frame) change and the generation of a premature stop codon, p.(Val417Alafs*76). A non-transfected HEK293T cell line was used as control.

Fig. 3 – Clinical appearance of female individuals with de novo variants in MED12.

(A) Females with truncating variants presented with marked facial dysmorphism such as abnormal skull shape, low-set, posteriorly rotated ears, blepharophimosis, crowded teeth, retrognathia and variable other abnormalities such as syndactyly of fingers and/or toes. Individuals 1 and 2 with a splice site variant share a very prominent forehead and MRI anomalies such as partial agenesis of corpus callosum, enlarged ventricles and tectum and cerebellar vermis abnormalities. (B) In comparison, females with *de novo* missense variants presented with minor, but rather non-specific facial dysmorphisms and less frequently with other structural or morphological abnormalities such as syndactyly and clinodactyly of fingers and/or toes (I12, I11, I16) or brain abnormalities such as agenesis of corpus callosum (I8, I15) or Chiari malformation (I18).

References

1. Jeronimo C, Robert F. The Mediator Complex: At the Nexus of RNA Polymerase II Transcription. *Trends Cell Biol.* 2017;27(10):765-783. doi:10.1016/j.tcb.2017.07.001
2. Conaway RC, Conaway JW. Function and regulation of the Mediator complex. *Curr Opin Genet Dev.* 2011;21(2):225-230. doi:10.1016/j.gde.2011.01.013
3. Poss ZC, Ebmeier CC, Taatjes DJ. The Mediator complex and transcription regulation. *Crit Rev Biochem Mol Biol.* 2013;48(6):575-608. doi:10.3109/10409238.2013.840259
4. Yin JW, Wang G. The Mediator complex: a master coordinator of transcription and cell lineage development. *Development.* 2014;141(5):977-987. doi:10.1242/dev.098392
5. Lyons MJ. MED12-Related Disorders. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*®. University of Washington, Seattle; 1993. Accessed April 30, 2020. <http://www.ncbi.nlm.nih.gov/books/NBK1676/>
6. Zhou H, Kim S, Ishii S, Boyer TG. Mediator modulates Gli3-dependent Sonic hedgehog signaling. *Mol Cell Biol.* 2006;26(23):8667-8682. doi:10.1128/MCB.00443-06
7. Kim S, Xu X, Hecht A, Boyer TG. Mediator is a transducer of Wnt/beta-catenin signaling. *J Biol Chem.* 2006;281(20):14066-14075. doi:10.1074/jbc.M602696200
8. Clark RD, Graham JM, Friez MJ, et al. FG syndrome, an X-linked multiple congenital anomaly syndrome: the clinical phenotype and an algorithm for diagnostic testing. *Genet Med.* 2009;11(11):769-775. doi:10.1097/GIM.0b013e3181bd3d90
9. Opitz JM, Kaveggia EG. Studies of malformation syndromes of man 33: the FG syndrome. An X-linked recessive syndrome of multiple congenital anomalies and mental retardation. *Z Kinderheilkd.* 1974;117(1):1-18.

10. Lujan JE, Carlin ME, Lubs HA. A form of X-linked mental retardation with marfanoid habitus. *Am J Med Genet.* 1984;17(1):311-322. doi:10.1002/ajmg.1320170124
11. Schwartz CE, Tarpey PS, Lubs HA, et al. The original Lujan syndrome family has a novel missense mutation (p.N1007S) in the MED12 gene. *Journal of medical genetics.* 2007;44(7):472-477. doi:10.1136/jmg.2006.048637
12. Maat-Kievit A, Brunner HG, Maaswinkel-Mooij P. Two additional cases of the Ohdo blepharophimosis syndrome. *Am J Med Genet.* 1993;47(6):901-906. doi:10.1002/ajmg.1320470618
13. Vulto-van Silfhout AT, de Vries BB, van Bon BW, et al. Mutations in MED12 cause X-linked Ohdo syndrome. *American journal of human genetics.* 2013;92(3):401-406. doi:10.1016/j.ajhg.2013.01.007
14. Bouazzi H, Lesca G, Trujillo C, Alwasayah MK, Munnich A. Nonsyndromic X-linked intellectual deficiency in three brothers with a novel MED12 missense mutation [c.5922G>T (p.Glu1974His)]. *Clin Case Rep.* 2015;3(7):604-609. doi:10.1002/ccr3.301
15. Lesca G, Moizard MP, Bussy G, et al. Clinical and neurocognitive characterization of a family with a novel MED12 gene frameshift mutation. *Am J Med Genet A.* 2013;161A(12):3063-3071. doi:10.1002/ajmg.a.36162
16. Rump P, Niessen RC, Verbruggen KT, Brouwer OF, de Raad M, Hordijk R. A novel mutation in MED12 causes FG syndrome (Opitz-Kaveggia syndrome). *Clin Genet.* 2011;79(2):183-188. doi:10.1111/j.1399-0004.2010.01449.x
17. Charzewska A, Maiwald R, Kahrizi K, et al. The power of the Mediator complex- Expanding the genetic architecture and phenotypic spectrum of MED12-related disorders. *Clin Genet.* 2018;94(5):450-456. doi:10.1111/cge.13412

18. Donnio LM, Bidon B, Hashimoto S, et al. MED12-related XLID disorders are dose-dependent of immediate early genes (IEGs) expression. *Hum Mol Genet.* 2017;26(11):2062-2075. doi:10.1093/hmg/ddx099
19. Hu H, Haas SA, Chelly J, et al. X-exome sequencing of 405 unresolved families identifies seven novel intellectual disability genes. *Mol Psychiatry.* 2016;21(1):133-148. doi:10.1038/mp.2014.193
20. Langley KG, Brown J, Gerber RJ, et al. Beyond Ohdo syndrome: A familial missense mutation broadens the MED12 spectrum. *Am J Med Genet A.* 2015;167A(12):3180-3185. doi:10.1002/ajmg.a.37354
21. Narayanan DL, Phadke SR. A novel variant in MED12 gene: Further delineation of phenotype. *Am J Med Genet A.* 2017;173(8):2257-2260. doi:10.1002/ajmg.a.38295
22. Niranjan TS, Skinner C, May M, et al. Affected kindred analysis of human X chromosome exomes to identify novel X-linked intellectual disability genes. *PLoS One.* 2015;10(2):e0116454. doi:10.1371/journal.pone.0116454
23. Patil SJ, Somashekar PH, Shukla A, et al. Clinical Variability in Familial X-Linked Ohdo Syndrome-Maat-Kievit-Brunner Type with MED12 Mutation. *J Pediatr Genet.* 2017;6(3):198-204. doi:10.1055/s-0037-1602386
24. Prescott TE, Kulseth MA, Heimdal KR, et al. Two male sibs with severe micrognathia and a missense variant in MED12. *Eur J Med Genet.* 2016;59(8):367-372. doi:10.1016/j.ejmg.2016.06.001
25. Prontera P, Ottaviani V, Rogaia D, et al. A novel MED12 mutation: Evidence for a fourth phenotype. *Am J Med Genet A.* 2016;170(9):2377-2382. doi:10.1002/ajmg.a.37805

26. Srivastava S, Niranjana T, May MM, et al. Dysregulations of sonic hedgehog signaling in MED12-related X-linked intellectual disability disorders. *Mol Genet Genomic Med*. 2019;7(4):e00569. doi:10.1002/mgg3.569
27. Tzschach A, Grasshoff U, Beck-Woedl S, et al. Next-generation sequencing in X-linked intellectual disability. *Eur J Hum Genet*. 2015;23(11):1513-1518. doi:10.1038/ejhg.2015.5
28. Yamamoto T, Shimojima K. A novel MED12 mutation associated with non-specific X-linked intellectual disability. *Hum Genome Var*. 2015;2:15018. doi:10.1038/hgv.2015.18
29. Murakami H, Enomoto Y, Tsurusaki Y, Sugio Y, Kurosawa K. A female patient with X-linked Ohdo syndrome of the Maat-Kievit-Brunner phenotype caused by a novel variant of MED12. *Congenit Anom (Kyoto)*. Published online July 19, 2019. doi:10.1111/cga.12350
30. Fieremans N, Van Esch H, Holvoet M, et al. Identification of Intellectual Disability Genes in Female Patients with a Skewed X-Inactivation Pattern. *Hum Mutat*. 2016;37(8):804-811. doi:10.1002/humu.23012
31. Wang C, Lin L, Xue Y, et al. MED12-Related Disease in a Chinese Girl: Clinical Characteristics and Underlying Mechanism. *Front Genet*. 2020;11:129. doi:10.3389/fgene.2020.00129
32. Firth HV, Richards SM, Bevan AP, et al. DECIPHER: Database of Chromosomal Imbalance and Phenotype in Humans Using Ensembl Resources. *Am J Hum Genet*. 2009;84(4):524-533. doi:10.1016/j.ajhg.2009.03.010
33. Sobreira N, Schiettecatte F, Valle D, Hamosh A. GeneMatcher: a matching tool for connecting investigators with an interest in the same gene. *Hum Mutat*. 2015;36(10):928-930. doi:10.1002/humu.22844

34. Allen RC, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW. Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *American journal of human genetics*. 1992;51(6):1229-1239.
35. Brogna S, Wen J. Nonsense-mediated mRNA decay (NMD) mechanisms. *Nat Struct Mol Biol*. 2009;16(2):107-113. doi:10.1038/nsmb.1550
36. Zweier C, Kraus C, Brueton L, et al. A new face of Borjeson-Forssman-Lehmann syndrome? De novo mutations in PHF6 in seven females with a distinct phenotype. *J Med Genet*. 2013;50(12):838-847. doi:10.1136/jmedgenet-2013-101918
37. Lyons MJ, Graham JM, Neri G, et al. Clinical experience in the evaluation of 30 patients with a prior diagnosis of FG syndrome. *J Med Genet*. 2009;46(1):9-13. doi:10.1136/jmg.2008.060509
38. Risheg H, Graham JM, Clark RD, et al. A recurrent mutation in MED12 leading to R961W causes Opitz-Kaveggia syndrome. *Nat Genet*. 2007;39(4):451-453. doi:10.1038/ng1992
39. Verloes A, Bremond-Gignac D, Isidor B, et al. Blepharophimosis-mental retardation (BMR) syndromes: A proposed clinical classification of the so-called Ohdo syndrome, and delineation of two new BMR syndromes, one X-linked and one autosomal recessive. *Am J Med Genet A*. 2006;140(12):1285-1296. doi:10.1002/ajmg.a.31270
40. Isidor B, Lefebvre T, Le Vaillant C, et al. Blepharophimosis, short humeri, developmental delay and hirschsprung disease: expanding the phenotypic spectrum of MED12 mutations. *Am J Med Genet A*. 2014;164A(7):1821-1825. doi:10.1002/ajmg.a.36539